arXiv:0809.1218v1 [math.DG] 7 Sep 2008

Contact Integrable Extensions of Symmetry Pseudo-Group and Coverings of the r-th Modified Dispersionless Kadomtsev – Petviashvili Equation

Oleg I. Morozov

Department of Mathematics, Moscow State Technical University of Civil Aviation,
Kronshtadtskiy Blvd 20, Moscow 125993, Russia
oim@foxcub.org

Abstract. We apply the technique of integrable extensions to the symmetry pseudo-group of the r-th mdKP equation. This gives another look on deriving known coverings and allows us to find new coverings for this equation.

AMS classification scheme numbers: 58H05, 58J70, 35A30

1. Introduction

Many useful methods to study nonlinear partial differential equations (PDEs) such as inverse scattering transformations, Bäcklund transformations, recursion operators, nonlocal symmetries and nonlocal conservation laws, can be successfully described in the framework of the theory of coverings, [22, 23, 24, 25]. Consequently, a problem to find a covering for a given PDE is of a significant importance. A number of techniques have been devised to handle this problem, [45, 12, 10, 18, 30, 31, 40, 19, 37, 38, 46, 43, 29, 16, 17]. In [28], examples of coverings of PDEs with three independent variables were found by means of Élie Cartan's method of equivalence, [5, 6, 7, 8, 15, 20, 39]. This approach was developed in [34, 35, 36], where it was shown that for a number of PDEs their coverings can be inferred from invariant linear combinations of Maurer–Cartan (MC) forms of the contact symmetry pseudo-groups of the PDEs. The aim of the present paper is to make the method of [34, 35, 36] more systematic. We apply the technique of integrable extensions of exterior differential systems, [4, §6], to MC forms of the symmetry pseudo-group of the r-th modified dispersionless Kadomtsev–Petviashvili equation, [1], and derive the known coverings, [14, 9, 21, 11, 36], and three new coverings.

2. Preliminaries

2.1. Coverings of PDEs

Let $\pi_\infty : J^\infty(\pi) \to \mathbb{R}^n$ be the infinite jet bundle of local sections of the bundle $\pi : \mathbb{R}^n \times \mathbb{R} \to \mathbb{R}$. The coordinates on $J^\infty(\pi)$ are (x^i, u_I), where $I = (i_1, ..., i_k)$ are symmetric multi-indices, $i_1, ..., i_k \in \{1, ..., n\}$, $u_\emptyset = u$, and for any local section f of π there exists a section $j_\infty(f) : \mathbb{R}^n \to J^\infty(\pi)$ such that $u_I(j_\infty(f)) = \partial^{\#I}(f)/\partial x^{i_1}...\partial x^{i_k}$, $\#I = \#(i_1, ..., i_k) = k$. The *total derivatives* on $J^\infty(\pi)$ are defined in the local coordintes as

$$D_i = \frac{\partial}{\partial x^i} + \sum_{\#I \ge 0} u_{Ii} \frac{\partial}{\partial u_I}.$$

We have $[D_i, D_j] = 0$ for $i, j \in \{1, ..., n\}$. A differential equation $F(x^i, u_K) = 0$ defines a submanifold $\mathcal{E}^\infty = \{D_I(F) = 0 \mid \#I \ge 0\} \subset J^\infty(\pi)$, where $D_I = D_{i_1} \circ ... \circ D_{i_k}$ for $I = (i_1, ..., i_k)$. We denote restrictions of D_i on $\mathcal{E}^\infty$ as $\bar{D}_i$.

In local coordinates, a *covering* over $\mathcal{E}^\infty$ is a bundle $\widetilde{\mathcal{E}}^\infty = \mathcal{E}^\infty \times \mathcal{V} \to \mathcal{E}^\infty$ with fibre coordinates v^α, $\alpha \in \{1, ..., N\}$ or $\alpha \in \mathbb{N}$, equipped with *extended total derivatives*

$$\widetilde{D}_i = \bar{D}_i + \sum_\alpha T_i^\alpha(x^j, u_I, v^\beta) \frac{\partial}{\partial v^\alpha}$$

such that $[\widetilde{D}_i, \widetilde{D}_j] = 0$ whenever $(x^i, u_I) \in \mathcal{E}^\infty$.

In terms of differential forms, the covering is defined by the forms, [45],

$$\omega^\alpha = dv^\alpha - T_i^\alpha(x^j, u_I, v^\beta)\, dx^i$$

such that

$$d\omega^\alpha \equiv 0 \mod \omega^\beta, \bar{\vartheta}_I \quad \Longleftrightarrow \quad (x^i, u_I) \in \mathcal{E}^\infty, \tag{1}$$

where $\bar{\vartheta}_I$ are restrictions of contact forms $\vartheta_I = du_I - u_{I,k}\,dx^k$ on $\mathcal{E}^\infty$.

2.2. Cartan's structure theory of Lie pseudo-groups

Let M be a manifold of dimension n. A *local diffeomorphism* on M is a diffeomorphism $\Phi : \mathcal{U} \to \widetilde{\mathcal{U}}$ of two open subsets of M. A *pseudo-group* $\mathfrak{G}$ on M is a collection of local diffeomorphisms of M, which is closed under composition *when defined*, contains an identity and is closed under inverse. A *Lie pseudo-group* is a pseudo-group whose diffeomorphisms are local analytic solutions of an involutive system of partial differential equations.

Élie Cartan's approach to Lie pseudo-groups is based on a possibility to characterize transformations from a pseudo-group in terms of a set of invariant differential 1-forms called *Maurer–Cartan forms*. In a general case MC forms ω^1, ..., ω^m of a Lie pseudo-group $\mathfrak{G}$ on M are defined on a direct product $\hat{M} \times G$, where $\mu : \hat{M} \times G \to M$ is a bundle, $m = \dim \hat{M}$, G is a finite-dimensional Lie group. The forms ω^i are semi-basic w.r.t. the natural projection $\hat{M} \times G \to \hat{M}$ and define a coframe on $\hat{M}$, that is, a basis of the cotangent bundle of $\hat{M}$. They characterize the pseudo-group $\mathfrak{G}$ in the following sense: a local diffeomorphism $\Phi : \mathcal{U} \to \widetilde{\mathcal{U}}$ on M belongs to $\mathfrak{G}$ whenever there exists a local diffeomorphism $\Psi : \mathcal{V} \to \widetilde{\mathcal{V}}$ on $\hat{M} \times G$ such that $\mu \circ \Psi = \Phi \circ \mu$ and the forms ω^j are invariant w.r.t. Ψ, that is,

$$\Psi^* \left(\omega^i|_{\widetilde{\mathcal{V}}}\right) = \omega^i|_{\mathcal{V}}. \tag{2}$$

Expressions of exterior differentials of the forms ω^i in terms of themselves give Cartan's structure equations of $\mathfrak{G}$:

$$d\omega^i = A^i_{\alpha j}\, \pi^\alpha \wedge \omega^j + B^i_{jk}\, \omega^j \wedge \omega^k, \qquad B^i_{jk} = -B^i_{kj}. \tag{3}$$

Here and below we assume summation on repeated indices. The forms π^α, $\alpha \in \{1, ..., \dim G\}$, are linear combinations of MC forms of the Lie group G and the forms ω^i. The coefficients $A^i_{\alpha j}$ and B^i_{jk} are either constants or functions of a set of invariants $U^\kappa : M \to \mathbb{R}$, $\kappa \in \{1, ..., l\}$, $l < \dim M$, of the pseudo-group $\mathfrak{G}$, so $\Phi^* \left(U^\kappa|_{\widetilde{\mathcal{U}}}\right) = U^\kappa|_{\mathcal{U}}$ for every $\Phi \in \mathfrak{G}$. In the latter case, differentials of U^κ are invariant 1-forms, so they are linear combinations of the forms ω^j,

$$dU^\kappa = C^\kappa_j\, \omega^j, \tag{4}$$

where the coefficients C^κ_j depend on the invariants U^1, ..., U^l only.

Eqs. (3) must be compatible in the following sense: we have

$$d(d\omega^i) = 0 = d\left(A^i_{\alpha j}\, \pi^\alpha \wedge \omega^j + B^i_{jk}\, \omega^j \wedge \omega^k\right), \tag{5}$$

therefore there must exist expressions

$$d\pi^\alpha = W^\alpha_{\lambda j}\, \chi^\lambda \wedge \omega^j + X^\alpha_{\beta\gamma}\, \pi^\beta \wedge \pi^\gamma + Y^\alpha_{\beta j}\, \pi^\beta \wedge \omega^j + Z^\alpha_{jk}\, \omega^j \wedge \omega^k, \tag{6}$$

with some additional 1-forms χ^λ and the coefficients $W^\alpha_{\lambda j}$ to Z^α_{jk} depending on the invariants U^κ such that the right-hand side of (5) appear to be identically equal to zero after substituting for (3), (4), and (6). Also, from (4) it follows that the right-hand side of the equation

$$d(dU^\kappa) = 0 = d(C^\kappa_j\,\omega^j) \tag{7}$$

must be identically equal to zero after substituting for (3) and (4).

The forms π^α are not invariant w.r.t. the pseudo-group $\mathfrak{G}$. Respectively, the structure equations (3) are not changing when replacing $\pi^\alpha \mapsto \pi^\alpha + z^\alpha_j\,\omega^j$ for certain parametric coefficients z^α_j. The dimension $r^{(1)}$ of the linear space of these coefficients satisfies the following inequality

$$r^{(1)} \le n\,\dim\,G - \sum_{k=1}^{n-1}(n-k)\,\sigma_k, \tag{8}$$

where the *reduced characters* σ_k are defined by

$$\sigma_k = \max_{u_1,...,u_k}\ \mathrm{rank}\ \mathbb{A}_k(u_1,...,u_k)$$

with the matrices $\mathbb{A}_k$ inductively defined by

$$\mathbb{A}_1(u_1) = \left(A^i_{\alpha j}\,u^j_1\right), \qquad \mathbb{A}_l(u_1,...,u_l) = \left(\begin{array}{c} \mathbb{A}_{l-1}(u_1,...,u_{l-1}) \\ A^i_{\alpha j}\,u^j_l \end{array}\right),$$

see [5, §5], [39, Def. 11.4] for the full discussion. The system of forms ω^k is *involutive* if (8) is an equality, [5, §6], [39, Def. 11.7].

Cartan's fundamental theorems, [5, §§16, 22–24], [8], [44, §§16, 19, 20, 25,26], [41, §§14.1–14.3], state that for a Lie pseudo-group there exists a set of MC forms whose structure equations satisfy the compatibility and involutivity conditions; conversely, if Eqs. (3), (4) meet the compatibility conditions (5), (7) and the involutivity condition, then there exists a collection of 1-forms ω^1, ... , ω^m and functions U^1, ... , U^l which satisfy (3) and (4). Eqs. (2) then define local diffeomorphisms from a Lie pseudo-group.

Example 1. Consider the bundle $J^2(\pi)$ of jets of the second order of the bundle π. A differential 1-form ϑ on $J^2(\pi)$ is called a *contact form* if it is annihilated by all 2-jets of local sections: $j_2(f)^*\vartheta = 0$. In the local coordinates every contact 1-form is a linear combination of the forms $\vartheta_0 = du - u_i\,dx^i$, $\vartheta_i = du_i - u_{ij}\,dx^j$, $i, j \in \{1, ..., n\}$, $u_{ji} = u_{ij}$. A local diffeomorphism $\Delta : J^2(\pi) \to J^2(\pi)$, $\Delta : (x^i, u, u_i, u_{ij}) \mapsto (\widetilde{x}^i, \widetilde{u}, \widetilde{u}_i, \widetilde{u}_{ij})$, is called a *contact transformation* if for every contact 1-form $\widetilde{\vartheta}$ the form $\Delta^*\widetilde{\vartheta}$ is also contact. We denote by $\mathrm{Cont}(J^2(\pi))$ the pseudo-group of contact transformations on $J^2(\pi)$. Consider the following 1-forms

$$\Theta_0 = a\,\vartheta_0, \quad \Theta_i = g_i\,\Theta_0 + a\,B^k_i\,\vartheta_k, \quad \Xi^i = c^i\,\Theta_0 + f^{ik}\,\Theta_k + b^i_k\,dx^k,$$
$$\Theta_{ij} = a\,B^k_i\,B^l_j\,(du_{kl} - u_{klm}\,dx^m) + s_{ij}\,\Theta_0 + w^k_{ij}\,\Theta_k, \tag{9}$$

defined on $J^2(\pi) \times \mathcal{H}$, where $\mathcal{H}$ is an open subset of $\mathbb{R}^{(2n+1)(n+3)(n+1)/3}$ with local coordinates $(a,\ b^i_k,\ c^i,\ f^{ik},\ g_i,\ s_{ij},\ w^k_{ij},\ u_{ijk})$, $i, j, k, \in \{1, ..., n\}$, such that $a \ne 0$,

$\det(b^i_k) \neq 0$, $f^{ik} = f^{ki}$, $s_{ij} = s_{ji}$, $w^k_{ij} = w^k_{ji}$, $u_{ijk} = u_{ikj} = u_{jik}$, while (B^i_k) is the inverse matrix for the matrix (b^i_k). As it is shown in [33], the forms (9) are MC forms for $\mathrm{Cont}(J^2(\pi))$, that is, a local diffeomorphism $\widehat{\Delta} : J^2(\pi) \times \mathcal{H} \to J^2(\pi) \times \mathcal{H}$ satisfies the conditions $\widehat{\Delta}^* \,\widetilde{\Theta}_0 = \Theta_0$, $\widehat{\Delta}^* \,\widetilde{\Theta}_i = \Theta_i$, $\widehat{\Delta}^* \,\widetilde{\Xi}^i = \Xi^i$, and $\widehat{\Delta}^* \,\widetilde{\Theta}_{ij} = \Theta_{ij}$ if and only if it is projectable on $J^2(\pi)$, and its projection $\Delta : J^2(\pi) \to J^2(\pi)$ is a contact transformation. The structure equations for $\mathrm{Cont}(J^2(\pi))$ have the form

$$\begin{aligned}
d\Theta_0 &= \Phi^0_0 \wedge \Theta_0 + \Xi^i \wedge \Theta_i,\\
d\Theta_i &= \Phi^0_i \wedge \Theta_0 + \Phi^k_i \wedge \Theta_k,\\
d\Xi^i &= \Phi^0_0 \wedge \Xi^i - \Phi^i_k \wedge \Xi^k + \Psi^{i0} \wedge \Theta_0 + \Psi^{ik} \wedge \Theta_k,\\
d\Theta_{ij} &= \Phi^k_i \wedge \Theta_{kj} + \Phi^k_j \wedge \Theta_{ki} - \Phi^0_0 \wedge \Theta_{ij} + \Upsilon^0_{ij} \wedge \Theta_0 + \Upsilon^k_{ij} \wedge \Theta_k + \Xi^k \wedge \Theta_{ijk},
\end{aligned}$$

where the additional forms Φ^0_0, Φ^0_i, Φ^k_i, Ψ^{i0}, Ψ^{ij}, Υ^0_{ij}, Υ^k_{ij}, and Θ_{ijk} depend on differentials of the coordinates of $\mathcal{H}$.

Example 2. Suppose $\mathcal{E}$ is a second-order differential equation in one dependent and n independent variables. We consider $\mathcal{E}$ as a submanifold in $J^2(\pi)$. Let $\mathrm{Cont}(\mathcal{E})$ be the group of contact symmetries for $\mathcal{E}$. It consists of all the contact transformations on $J^2(\pi)$ mapping $\mathcal{E}$ to itself. Let $\iota_0 : \mathcal{E} \to J^2(\pi)$ be an embedding, and $\iota = \iota_0 \times \mathrm{id} : \mathcal{E} \times \mathcal{H} \to \mathrm{J}^2(\pi) \times \mathcal{H}$. The MC forms of $\mathrm{Cont}(\mathcal{E})$ can be computed from the forms $\theta_0 = \iota^*\Theta_0$, $\theta_i = \iota^*\Theta_i$, $\xi^i = \iota^*\Xi^i$, and $\theta_{ij} = \iota^*\Theta_{ij}$ by means of Cartan's method of equivalence, [5, 6, 7, 8, 15, 20, 39], see details and examples in [13, 32, 33].

3. Cartan's structure of the contact symmetry pseudo-group for r-mdKP

The r-th mdKP

$$u_{tx} = -\frac{(3-r)\,(1-r)}{2}\,u_x^2 u_{xx} + \frac{r\,(3-r)}{2-r}\,u_x u_{xy} + \frac{3-r}{(2-r)^2}\,u_{yy} + \frac{(3-r)\,(1-r)}{2-r}\,u_y u_{xx},$$

$r \in \mathbb{Z}\backslash\{2\}$, was derived in [1]. For a convenience of computations we use the following change of variables:

$$\tilde{t} = (3-r)\,t, \quad \tilde{x} = x, \quad \tilde{y} = (2-r)\,y, \quad \tilde{u} = -(1-r)\,u,$$

where $r \notin \{1,2,3\}$. Then we have

$$\tilde{u}_{\tilde{y}\tilde{y}} = \tilde{u}_{\tilde{t}\tilde{x}} + \left(\frac{1}{2\,(1-r)}\,\tilde{u}_{\tilde{x}}^2 + \tilde{u}_{\tilde{y}}\right)\tilde{u}_{\tilde{x}\tilde{x}} + \frac{r}{1-r}\,\tilde{u}_{\tilde{x}}\,\tilde{u}_{\tilde{x}\tilde{y}}.$$

We drop tildes and denote $\kappa = \frac{r}{1-r}$; this yields

$$u_{yy} = u_{tx} + \left(\tfrac{1}{2}\,(\kappa+1)\,u_x^2 + u_y\right)\,u_{xx} + \kappa\,u_x\,u_{xy}. \tag{10}$$

The exceptional cases $r = 2$ and $r = 3$ correspond to $\kappa = -2$ and $\kappa = -\frac{3}{2}$, respectively. We will not consider the case of $r = 1$. The case of $\kappa = -1$ is exceptional, too, since $r \to \infty$ when $\kappa \to -1$. In the cases of $\kappa = 0$, $\kappa = 1$, and $\kappa = -1$ Eq. (10) gets the forms of the mdKP equation, [28, 26, 27], the dBKP equation, [42, 21], and the equation describing Lorentzian hyper-CR Einstein–Weil structures, [11, 14].

We use the method outlined in the previous section to compute MC forms and structure equations of the pseudo-group of contact symmetries for Eq. (10). In the next section, the analysis of integrable extensions will depend crucially on invariants of the pseudo-group. So in the computations we shall not suppress the invariants, as it was done in [36]. We write out here the structure equations for the forms θ_0, θ_j, ξ^j with $j \in \{1, 2, 3\}$ only; the full sets of the structure equations are given in Appendix.

When $\kappa \neq -1$, we have

$$
\begin{aligned}
d\theta_0 &= \eta_1 \wedge \theta_0 + \xi^1 \wedge \theta_1 + \xi^2 \wedge \theta_2 + \xi^3 \wedge \theta_3,\\
d\theta_1 &= \tfrac{1}{16}\left(24\,\eta_1 - 24\,\theta_{22} - 12\,V\,\xi^1 - 12\,\xi^2 - (12\,U - \kappa + 4)\,\xi^3\right) \wedge \theta_1 + \xi^1 \wedge \theta_{11} + \xi^2 \wedge \theta_{12}\\
&\quad + \left(\tfrac{1}{8}\,(2\,\kappa^2 + 15\,\kappa + 4)\,\theta_2 + \kappa\,\theta_{23} + (\kappa\,U + 1 - \tfrac{1}{8}\,\kappa\,(\kappa - 10))\,\xi^2\right.\\
&\quad + \left.\tfrac{1}{16}\,(7\,\kappa + 4)\,V\,\xi^3\right) \wedge \theta_0 + ((\kappa + 1)\,\theta_2 + (\kappa + 2)\,\xi^2) \wedge \theta_3 + \xi^3 \wedge \theta_{13},\\
d\theta_2 &= \tfrac{1}{2}\left(\eta_1 - \theta_{22} - \tfrac{1}{2}\,V\,\xi^1 - \xi^2 - \left(U - \tfrac{7}{8}\,\kappa - \tfrac{1}{2}\right)\,\xi^3\right) \wedge \theta_2 + \xi^1 \wedge \theta_{12} + \xi^2 \wedge \theta_{22} + \xi^3 \wedge \theta_{23},\\
d\theta_3 &= \left(\eta_1 - \theta_{22} - \tfrac{1}{2}\,V\,\xi^1 - \xi^2 - \left(U - \tfrac{1}{4}\,\kappa\right)\,\xi^3\right) \wedge \theta_3 + \tfrac{1}{2}\,(\kappa + 2)\,\xi^2 \wedge \theta_2 + \xi^1 \wedge \theta_{13}\\
&\quad + \tfrac{1}{8}\,(\kappa - 4)\,\left(\theta_{22} + \xi^2 + \tfrac{1}{8}\,(8\,U - \kappa - 4)\,\xi^3\right) \wedge \theta_0 + \xi^2 \wedge \theta_{23} + \xi^3 \wedge \theta_{12},\\
d\xi^1 &= -\tfrac{1}{2}\,\left(\eta_1 - 3\,\theta_{22} + 3\,\xi^2 - \left(3\,U - \tfrac{1}{8}\,(\kappa - 4)\right)\,\xi^3\right) \wedge \xi^1,\\
d\xi^2 &= \tfrac{1}{2}\,\left(\eta_1 + \theta_{22} + \tfrac{1}{2}\,V\,\xi^1 + \left(U + \tfrac{1}{8}\,\kappa + \tfrac{3}{2}\right)\,\xi^3\right) \wedge \xi^2 + \left(\tfrac{1}{8}\,(\kappa - 4)\,\theta_0 - \theta_3\right) \wedge \xi^1 - \theta_2 \wedge \xi^3,\\
d\xi^3 &= \left(\theta_{22} + \tfrac{1}{2}\,V\,\xi^1 + \xi^2\right) \wedge \xi^3 - (\kappa + 2)\,\left(\theta_2 + \xi^2\right) \wedge \xi^1,
\end{aligned}
$$

where the invariants

$$
U = \frac{u_{xxy} + u_x\,u_{xxx}}{u_{xx}^2}, \tag{11}
$$

$$
V = \frac{u_{xxx}\,(2\,u_{txx} + ((\kappa + 3)\,u_x^2 + 2\,u_y)\,u_{xxx} + 2\,(\kappa + 2)\,(u_x\,(u_{xxy} + u_{xx}^2) + u_{xx}\,u_{xy}))}{u_{xx}^4}, \tag{12}
$$

satisfy

$$
\begin{aligned}
dU &= \theta_2 - U\,\theta_{22} - \eta_2 - \tfrac{1}{64}\,V\,\left(64\,U + 7\,\kappa^2 + 38\,\kappa - 8\right)\,\xi^1 - \tfrac{1}{16}\,\left(8\,U - \kappa - 6\right)\,\xi^2\\
&\quad - \left(2\,U^2 - (2\,\kappa + 1)\,U - \tfrac{3}{4}\,V - \kappa - 1\right)\,\xi^3,\\
dV &= -\tfrac{1}{4}\,(\kappa - 4)\,\theta_0 + \tfrac{1}{4}\,\left(8\,(\kappa + 2)\,U + 3\,(\kappa^2 + 6\,\kappa + 8)\right)\,\theta_2 + 2\,\theta_3 - \tfrac{3}{2}\,V\,\theta_{22}\\
&\quad + 2\,(\kappa + 2)\,\theta_{23} - 2\,\eta_3 + \tfrac{1}{2}\,V\,\eta_1 + (2\,(\kappa + 2)\,(2\,U + 1) - V)\,\xi^2\\
&\quad - \tfrac{1}{32}\,V\,\left(80\,U + 7\,\kappa^2 + 4\,\kappa - 64\right)\,\xi^3,
\end{aligned}
$$

and where

$$
\begin{aligned}
\theta_0 &= \frac{u_{xxx}^2}{u_{xx}^3}\,\vartheta_0,\\
\theta_1 &= \frac{u_{xxx}^3}{{u_{xx}}^6}\,\left(\kappa\,(u_x\,u_{xx} + u_{xy})\,\vartheta_0 + \vartheta_1 + \left(\tfrac{1}{2}\,(\kappa + 3)\,{u_x}^2 + u_y\right)\,\vartheta_2 + (\kappa + 2)\,u_x\,\vartheta_3\right),\\
\theta_2 &= \frac{u_{xxx}}{u_{xx}^2}\,\vartheta_2,\\
\theta_3 &= \frac{u_{xxx}^2}{u_{xx}^4}\,\left((\kappa - 4)\,u_{xx}\,\vartheta_0 + u_x\,\vartheta_2 + \vartheta_3\right),\\
\theta_{22} &= \frac{1}{u_{xx}}\,\vartheta_{22},
\end{aligned}
\tag{13}
$$

$$
\begin{aligned}
\xi^1 &= \frac{u_{xx}^3}{u_{xxx}}\,dt,\\
\xi^2 &= \frac{u_{xxx}}{u_{xx}}\,\left(\left(\tfrac{1}{2}\,(\kappa+1)\,u_{xx}^2 - u_y\right)\,dt + dx - u_x\,dy\right),\\
\xi^3 &= -\,u_{xx}\,u_x\,(\kappa+2)\,dt + u_{xx}\,dy,\\
\eta_1 &= 2\,\frac{du_{xxx}}{u_{xxx}} - 3\,(\theta_{22}-\xi^2) - \frac{1}{2\,u_{xx}^3}\,\left(3\,V\,u_{xx}^3 - 4\,(\kappa+3)\,(u_x\,u_{xx}+u_{xy})\,u_{xxx}\right)\,\xi^1\\
&\quad -\,\left(3\,U + \tfrac{1}{8}(\kappa-4)\right)\,\xi^3,
\end{aligned}
$$

with

$$
\begin{aligned}
\vartheta_0 &= du - u_t\,dt - u_x\,dx - u_y\,dy,\\
\vartheta_1 &= du_t - u_{tt}\,dt - u_{tx}\,dx - u_{ty}\,dy,\\
\vartheta_2 &= du_x - u_{tx}\,dt - u_{xx}\,dx - u_{xy}\,dy,\\
\vartheta_3 &= du_y - u_{ty}\,dt - u_{xy}\,dx - (u_{tx} + \left(\tfrac{1}{2}(\kappa+1)\,u_x^2 + u_y\right)\,u_{xx} + \kappa\,u_x\,u_{xy})\,dy,\\
\vartheta_{22} &= du_{xx} - u_{txx}\,dt - u_{xxx}\,dx - u_{xxy}\,dy.
\end{aligned}
$$

We need not explicit expressions for the other MC forms in the sequel.

For $\kappa = -1$, we get

$$
\begin{aligned}
d\theta_0 &= \eta_1\wedge\theta_0 + \xi^1\wedge\theta_1 + \xi^2\wedge\theta_2 + \xi^3\wedge\theta_3,\\
d\theta_1 &= \tfrac{3}{2}\,\eta_1\wedge\theta_1 + 2\,\eta_2\wedge\theta_3 + \eta_3\wedge\theta_0 - \tfrac{1}{8}\,\left(12\,\theta_{22} + 16\,U\,\xi^1 + 12\,\xi^2 + 7\,\xi^3\right)\wedge\theta_1 + \xi^1\wedge\theta_{11}\\
&\quad +\,\xi^2\wedge\theta_{12} + \xi^3\wedge\theta_{13},\\
d\theta_2 &= \tfrac{1}{8}\,\left(4\,(\eta_1 - \theta_{22} - \xi^2) - 3\,\xi^3\right)\wedge\theta_2 + \xi^1\wedge\theta_{12} + \xi^2\wedge\theta_{22} + \xi^3\wedge\theta_{23},\\
d\theta_3 &= \left(\eta_1 - \theta_{22} - U\,\xi^1 - \xi^2 - \tfrac{5}{8}\,\xi^3\right)\wedge\theta_3 + \eta_2\wedge\theta_2 - \tfrac{5}{8}\,(\theta_{22}+\xi^2)\wedge\theta_0 + \xi^1\wedge\theta_{13}\\
&\quad +\,\xi^2\wedge\theta_{23} + \xi^3\wedge\theta_{12},\\
d\xi^1 &= \tfrac{1}{8}\,\left(12\,\theta_{22} - 4\,\eta_1 + 12\,\xi^2 + 7\,\xi^3\right)\wedge\xi^1,\\
d\xi^2 &= -\,\tfrac{1}{8}\,(5\,\theta_0 + 8\,\theta_3)\wedge\xi^1 + \tfrac{1}{8}\,\left(4\,(\eta_1+\theta_{22}) + 3\,\xi^3\right)\wedge\xi^2 - (\eta_2+\theta_2)\wedge\xi^3,\\
d\xi^3 &= -\,(2\,\eta_2 + \theta_2)\wedge\xi^1 + (\theta_{22} + U\,\xi^1 + \xi^2)\wedge\xi^3,
\end{aligned}
$$

where the invariant

$$
U = \frac{u_{xxx}\,(u_x\,u_{xxy} - u_{txx} - u_y\,u_{xxx} - u_{xy}\,u_{xx}) + u_{xxy}\,(u_{xxy} + u_{xx}^2)}{u_{xx}^4} - \frac{33}{64} \tag{14}
$$

satisfies

$$
dU = -\tfrac{5}{8}\,\theta_0 + \tfrac{9}{8}\,\theta_2 - \tfrac{1}{8}\,(4\,U-7)\,\xi^2 - \tfrac{3}{2}\,U\,(\theta_{22}+\xi^3) + \theta_{23} + \eta_3 - \tfrac{1}{4}\,\eta_2 + \tfrac{1}{2}\,U\,\eta_1, \tag{15}
$$

and where

$$
\begin{aligned}
\theta_0 &= \frac{u_{xxx}^2}{{u_{xx}}^3}\,\vartheta_0,\\
\theta_1 &= \frac{u_{xxx}^3}{u_{xx}^6}\,\vartheta_1 + \frac{u_{xxx}}{64\,u_{xx}^6}\,\left(64\,(u_y\,u_{xxx}^2 + u_{xxy}^2) + u_{xx}^2\,(9\,u_{xx}^2 - 48\,u_{xxy})\right)\,\vartheta_2\\
&\quad -\,\frac{u_{xxx}^2}{u_{xx}^7}\,\left(u_{xxx}\,(u_{txx} + u_y\,u_{xxx} - u_x\,u_{xxy} + 2\,u_{xx}\,u_{xy}) - u_{xxy}\,(u_{xxy} + 2\;u_{xx}^2) + \tfrac{57}{64}\,u_{xx}^4\right)\,\vartheta_0
\end{aligned}
$$

$$\begin{aligned}
&\quad - \frac{u_{xxx}^2}{4\,u_{xx}^6}\,\left(8\,u_{xxy}+4\,u_x\,u_{xxx}-3\,u_{xx}^2\right)\,\vartheta_3,\\
\theta_2 &= \frac{u_{xxx}}{u_{xx}^2}\,\vartheta_2,\\
\theta_3 &= \frac{u_{xxx}}{8u_{xx}^4}\,\left(8\,u_{xxx}\,\vartheta_3-5\,u_{xx}\,u_{xxx}\,\vartheta_0-(8\,u_{xxy}-3\,u_{xx}^2)\vartheta_2\right),\\
\theta_{22} &= \frac{1}{u_{xx}}\,\vartheta_{22},\\
\xi^1 &= \frac{u_{xx}^3}{u_{xxx}}\,dt,\\
\xi^2 &= \frac{1}{u_{xx}u_{xxx}}\,\left(u_{xxy}\,\left(u_{xxy}-\tfrac{3}{4}\,u_{xx}^2\right)+u_{xxx}\,(u_x\,u_{xxy}-u_y\,u_{xxx})\right)\,dt\\
&\quad -\tfrac{3}{64}\,u_{xx}^2\,\left(8\,u_x\,u_{xxx}-3\,u_{xx}^2\right)+\frac{u_{xxx}}{u_{xx}}\,dx+\frac{(8\,u_{xxy}-3\,u_{xx}^2)}{8\,u_{xx}}\,dy,\\
\xi^3 &= \frac{u_{xx}}{4u_{xxx}}\,\left(8\,u_{xxy}+4\,u_x\,u_{xxx}-3\,u_{xx}^2\right)\,dt+u_{xx}\,dy,\\
\eta_1 &= 2\,\frac{du_{xxx}}{u_{xxx}}-3\,(\theta_{22}+\xi^2)+\frac{1}{2u_{xx}^3}\,\left((4\,U+3)\,u_{xx}^3\,+8\,(u_{xy}\,u_{xxx}-u_{xx}\,u_{xxy})\right)\,\xi^1-\tfrac{1}{2}\,\xi^3.
\end{aligned} \tag{16}$$

4. Integrable extensions

In [4, §6], the definition of integrable extension of an exterior differential system is designed to study finite-dimensional coverings. In general case, coverings of PDEs with three or more independent variables are infinite-dimensional, [29]. To cope with infinite-dimensional coverings we use a natural generalization of the definition.

Suppose $\mathfrak{G}$ is a Lie pseudo-group on a manifold M and ω^1, ... , ω^m are its MC forms with structure equations (3), (4). Consider a system of equations

$$\begin{aligned}
d\tau^q &= D^q_{\rho r}\,\eta^\rho\wedge\tau^r+E^q_{rs}\,\tau^r\wedge\tau^s+F^q_{r\beta}\,\tau^r\wedge\pi^\beta+G^q_{rj}\,\tau^r\wedge\omega^j+H^q_{\beta j}\,\pi^\beta\wedge\omega^j\\
&\quad + I^q_{jk}\,\omega^j\wedge\omega^k,
\end{aligned} \tag{17}$$

$$dV^\epsilon = J^\epsilon_j\,\omega^j+K^\epsilon_q\,\tau^q, \tag{18}$$

with unknown 1-forms τ^q, $q\in\{1,...,Q\}$, η^ρ, $\rho\in\{1,...,R\}$, and unknown functions V^ϵ, $\epsilon\in\{1,...,S\}$ for some $Q,R,S\in\mathbb{N}$. The coefficients $D^q_{\rho r}$, ..., K^ϵ_q in (17), (18) are supposed to be functions of U^q and V^γ.

Definition 1. System (17), (18) is an *integrable extension* of system (3), (4), if Eqs. (17), (18), (3), and (4) together satisfy the compatibility and involutivity conditions.

In this case from Cartan's third fundamental theorem for Lie pseudo-groups it follows that there exists a set of forms τ^q and functions V^ϵ which are solutions to Eqs. (17) and (18). Then τ^q, V^ϵ together with ω^i, U^q define a Lie pseudo-group on a manifold $N\cong M\times\mathbb{R}^Q$.

Definition 2. The integrable extension is called *trivial*, if there is a change of variables on N such that in the new variables the coefficients $F^q_{r\beta}$, G^q_{rj}, $H^q_{\beta j}$, I^q_{jk}, and J^ϵ_j are equal to zero, while the coefficients $D^q_{\rho r}$, E^q_{rs}, and K^ϵ_q are independent of U^q. Otherwise, the integrable extension is called *non-trivial.*

Let θ^α_I and ξ^j be a set of MC forms of the symmetry pseudo-group $\mathrm{Cont}(\mathcal{E})$ of a PDE $\mathcal{E}$ such that $\xi^1 \wedge ... \wedge \xi^n \neq 0$ on any solution manifold of $\mathcal{E}$, while θ^α_I are contact forms. We take the following reformulation of the definition (1) of a covering.

Definition 3. A non-trivial integrable extension of the form

$$d\omega^q = \Pi^q_r \wedge \omega^r + \xi^j \wedge \Omega^q_j \tag{19}$$

is referred to as a *contact integrable extension* (CIE) of the structure equations of $\mathrm{Cont}(\mathcal{E})$ if

(1) Π^q_r are some non-trivial differential 1-forms,
(2) $\Omega^q_j \equiv 0 \mod \theta^\alpha_I, \omega^q_j$ for some additional 1-forms ω^q_j.

Since (19) is integrable extension, Cartan's theorem yields existence of the forms ω^q satisfying (19). From [3, Ch. IV, Prop. 5.10] it follows that the forms ω^q define a system of PDEs. This system is a covering for $\mathcal{E}$.

We apply this construction to the structure equations (41) and (42) of the symmetry pseudo-group of the r-th mdKP equation. We restrict our analysis to CIEs of the form

$$\begin{aligned} d\omega_0 = {} & \left(\sum_{i=0}^{3} A_i\,\theta_i + \sum{}^{*} B_{ij}\,\theta_{ij} + \sum_{s=1}^{7} C_s\,\eta_s + \sum_{j=1}^{3} D_j\,\xi^j + E\,\omega_1\right) \wedge \omega_0 \\ & + \sum_{j=1}^{3}\left(\sum_{k=0}^{3} F_{jk}\,\theta_k + G_j\,\omega_1\right) \wedge \xi^j, \end{aligned} \tag{20}$$

where $\sum{}^{*}$ means summation for all $i, j \in \mathbb{N}$ such that $1 \le i \le j \le 3$, $(i, j) \neq (3, 3)$, and consider two cases:

1) the coefficients A_i to G_j in (20) depend on the invariants of the symmetry pseudo-group of Eq. (10) only, i.e., on the U and V when $\kappa \neq -1$ and on U when $\kappa = -1$;

2) the coefficients A_i to G_j depend also on one additional invariant, say W. In this case, the differential of this new invariant satisfies the following equation

$$dW = \sum_{i=0}^{3} H_i\,\theta_i + \sum{}^{*} I_{ij}\,\theta_{ij} + \sum_{s=1}^{7} J_s\,\eta_s + \sum_{j=1}^{3} K_j\,\xi^j + \sum_{q=0}^{1} L_q\,\omega_q, \tag{21}$$

where the coefficients H_i to L_q are functions of U, V, W when $\kappa \neq -1$ or of U and W when $\kappa = -1$.

The requirements of Defintions 1 and 3 yield over-determined systems for the coefficients A_i to G_j in the first case and A_i to L_q in the second case. The results of analysis of these systems are summarized in the following theorems.

Theorem 1. *Let $\kappa \neq -1$ and suppose that the coefficients A_i to G_j in (20) depend on the invariants U, V defined by (11) and (12). Then* CIE *(20) of (41) is contact-equivalent* *to the following equation:*

$$\begin{aligned} d\omega_0 = &\left(\omega_1 + \tfrac{1}{2}\,(\eta_1 + \theta_{22}) + \tfrac{1}{4}\,V\,\xi^1 + \tfrac{1}{16}\,(8\,U + \kappa + 12)\,\xi^3\right) \wedge \omega_0 + \omega_1 \wedge \xi^2 + \theta_2 \wedge \xi^3 \\ &- \left(\tfrac{1}{8}\,(\kappa - 4)\,\theta_0 - \theta_3\right) \wedge \xi^1. \end{aligned} \tag{22}$$

Theorem 2. *For $\kappa \notin \{-3, -1\}$ suppose that the coefficients A_i to G_j in (20) and H_i to L_q in (21) depend on the invariants U, V, and W, where U and V are defined by (11), (12). Then* CIE *(20), (21) of (41) is contact-equivalent to the following equations:*

$$\begin{aligned} d\omega_0 = &\left(\omega_1 + \tfrac{1}{2}\,(\eta_1 + \theta_{22}) + \tfrac{1}{4}\,\left(V - 2\,W\,(W-2)\right)\xi^1 + \tfrac{1}{16}\,(8\,(U - W) + \kappa + 12)\,\xi^3\right) \wedge \omega_0 \\ &+ \left(W^2\,\omega_1 - \tfrac{1}{8}\,(\kappa - 4)\,\theta_0 + W\,\theta_2 + \theta_3\right) \wedge \xi^1 + \omega_1 \wedge \xi^2 + (W\,\omega_1 + \theta_2) \wedge \xi^3, \end{aligned} \tag{23}$$

$$\begin{aligned} dW = &\tfrac{1}{16}\,W\,\left(8\,(\eta_1 - \theta_{22} - \kappa\,\omega_1 - (\kappa+1)\,\omega_2) - 4\,\left(V - 2\,W\,(W - 2\,(\kappa + 1))\right)\,\xi^1\right. \\ &- \left.(8\,(U - W) + 5\,(3\kappa + 4))\,\xi^3\right). \end{aligned} \tag{24}$$

In the case of $\kappa = -3$, every CIE *(20), (21) of (41) is contact-equivalent either to (23), (24) or to the following equations:*

$$\begin{aligned} d\omega_0 = &\left(\omega_1 + \tfrac{1}{2}\,(\eta_1 + \theta_{22}) + \tfrac{1}{4}\,V\,\xi^1 + \tfrac{3}{2}\,\xi^2 + \tfrac{1}{16}\,(8\,(U + 2\,W) + 9)\,\xi^3\right) \wedge \omega_0 \\ &+ \left(\left(W + \tfrac{7}{8}\right)\,\theta_0 + \theta_3\right) \wedge \xi^1 + \omega_1 \wedge \xi^2 + \theta_2 \wedge \xi^3, \end{aligned} \tag{25}$$

$$dW = W\,\left(\omega_1 - \theta_{22} - \tfrac{1}{2}\,V\,W\,\xi^1 - (U - W - 1)\,\xi^3\right). \tag{26}$$

Theorem 3. *In the case of $\kappa = -1$ there are no* CIE*s of (42) with the coefficients A_i to G_j depending on the invariant U from (14) only. Every* CIE *(20), (21) of (42) whose coefficients A_i to L_q depend on U and W is contact-equivalent to the following equations:*

$$\begin{aligned} d\omega_0 = &\left(\omega_1 + \tfrac{1}{2}\,(\eta_1 + \theta_{22} - W\,(W-2)\,\xi^1) - \tfrac{1}{8}\,(4\,W - 7)\,\xi^3\right) \wedge \omega_0 \\ &+ \left(W^2\,\omega_1 + \tfrac{5}{8}\,\theta_0 + W\,\theta_2 + \theta_3\right) \wedge \xi^1 + \omega_1 \wedge \xi^2 + (W\,\omega_1 + \theta_2) \wedge \xi^3, \end{aligned} \tag{27}$$

$$dW = \eta_2 + U\,W\,\xi^1 + \tfrac{1}{2}\,W\,\left(\eta_1 - \theta_{22} - (W + 1)\,\xi^2 + (2\,U - W)\,\xi^3\right). \tag{28}$$

Since forms (13) in (22) – (26) and forms (16) in (27), (28) are known, it is easy to find the forms ω_0 explicitly:

Theorem 4. *We have the following solutions to Eqs. (22), (23), (25), and (27) up to a contact equivalence:*

1) Eq. (22) yields

$$\omega_0 = \frac{u_{xxx}}{u_{xx}\,v_x}\,\left(dv - \left(\tfrac{1}{2}\,(\kappa + 1)\,u_x^2 - u_y\right)\,v_x\,dt - v_x\,dx + u_x\,v_x\,dy\right). \tag{29}$$

2) Integrating Eqs. (23), (24) yields the following results:

when $\kappa \notin \{-2, -\frac{3}{2}, -1\}$, we have

$$\begin{aligned} \omega_0 = &\frac{u_{xxx}}{u_{xx}\,v_x}\,\left(dv - \left((2\kappa + 3)^{-1}\,v_x^{2\,(\kappa+1)} - \lambda\,u_x\,v_x^{\kappa+1} + \left(\tfrac{1}{2}\,(\kappa + 1)\,u_x^2 - \,u_y\right)\right)\,v_x\,dt\right. \\ &- \left. v_x\,dx - \left((\kappa + 2)^{-1}\,v_x^{\kappa+1} - u_x\right)\,v_x\,dy\right), \end{aligned} \tag{30}$$

when $\kappa = -2$, we have

$$\omega_0 = \frac{u_{xxx}}{u_{xx}v_x}\,\left(dv + \left(v_x^{-1} + u_x + \left(\tfrac{1}{2}\,u_x^2 + u_y\right)\,v_x\right)\,dt - \left(\ln|v_x| - u_x\,v_x\right)\,dy - v_x\,dx\right), \tag{31}$$

and when $\kappa = -\frac{3}{2}$, we have

$$\omega_0 = \frac{u_{xxx}}{u_{xx}\, v_x}\, \Big(dv + \Big(\big(\tfrac{1}{4}\, u_x^2 + u_y\big)\; v_x + u_x \sqrt{|v_x|} - \ln |v_x| \Big)\; dt + \Big(u_x\, v_x - 2\, \sqrt{|v_x|} \Big)\; dy$$
$$-\;\; v_x\, dx\big)\,. \tag{32}$$

3) From Eqs. (25), (26) we get

$$\omega_0 = \frac{u_{xxx}}{u_{xx}\, v_x}\, \big(dv + \big(u + (u_x^2 + u_y)\, v_x \big)\; dt - v_x\, dx + (v_x\, u_x + x)\, dy \big)\,. \tag{33}$$

4) Eqs. (27), (28) give

$$\omega_0 = \frac{u_{xxx}}{u_{xx}\, v_x}\, \big(dv + v_x \big(u_y + \lambda\, u_x - \lambda^2 \big)\; dt - v_x\, dx + v_x\, (u_x - \lambda)\, dy \big)\,, \quad \lambda \in \mathbb{R} \tag{34}$$

Forms (29) – (34) define infinite-dimensional coverings of Eq. (10) with the fibre variables v_j, $j \in \mathbb{N} \cup \{0\}$, and the following extended total derivatives, respectively:

$$\left\{ \begin{array}{lcl} \widetilde{D}_t & = & \bar{D}_t + \sum\limits_{j=0}^{\infty} \widetilde{D}_x^j \left(\left(\frac{1}{2}\, (\kappa+1)\, u_x^2 - u_y \right)\; v_1 \right)\; \frac{\partial}{\partial v_j}, \\ \widetilde{D}_x & = & \bar{D}_x + \sum\limits_{j=0}^{\infty} v_{j+1}\, \frac{\partial}{\partial v_j}, \\ \widetilde{D}_y & = & \bar{D}_y - \sum\limits_{j=0}^{\infty} \widetilde{D}_x^j\, (u_x\, v_1)\; \frac{\partial}{\partial v_j}, \end{array} \right. \tag{35}$$

$$\left\{ \begin{array}{lcl} \widetilde{D}_t & = & \bar{D}_t + \sum\limits_{j=0}^{\infty} \widetilde{D}_x^j \left((2\kappa+3)^{-1}\, v_1^{2\,\kappa+3} - u_x\, v_1^{\kappa+2} + \left(\frac{1}{2}\, (\kappa+1)\, u_x^2 -\, u_y \right)\; v_1 \right)\; \frac{\partial}{\partial v_j}, \\ \widetilde{D}_x & = & \bar{D}_x + \sum\limits_{j=0}^{\infty} v_{j+1}\, \frac{\partial}{\partial v_j}, \\ \widetilde{D}_y & = & \bar{D}_y + \sum\limits_{j=0}^{\infty} \widetilde{D}_x^j \left((\kappa+2)^{-1}\, v_1^{\kappa+2} - u_x\, v_1 \right)\, \frac{\partial}{\partial v_j}, \end{array} \right. \tag{36}$$

$$\left\{ \begin{array}{lcl} \widetilde{D}_t & = & \bar{D}_t - \sum\limits_{j=0}^{\infty} \widetilde{D}_x^j \left(v_1^{-1} + u_x + \left(\frac{1}{2}\, u_x^2 + u_y \right)\; v_1 \right)\; \frac{\partial}{\partial v_j}, \\ \widetilde{D}_x & = & \bar{D}_x + \sum\limits_{j=0}^{\infty} v_{j+1}\, \frac{\partial}{\partial v_j}, \\ \widetilde{D}_y & = & \bar{D}_y + \sum\limits_{j=0}^{\infty} \widetilde{D}_x^j\, (\ln |v_1| - u_x\, v_1)\; \frac{\partial}{\partial v_j}, \end{array} \right. \tag{37}$$

$$\left\{ \begin{array}{lcl} \widetilde{D}_t & = & \bar{D}_t - \sum\limits_{j=0}^{\infty} \widetilde{D}_x^j \Big(\left(\frac{1}{4}\, u_x^2 + u_y \right)\; v_1 + u_x \sqrt{|v_1|} - \ln |v_1| \Big)\; \frac{\partial}{\partial v_j}, \\ \widetilde{D}_x & = & \bar{D}_x + \sum\limits_{j=0}^{\infty} v_{j+1}\, \frac{\partial}{\partial v_j}, \\ \widetilde{D}_y & = & \bar{D}_y + \sum\limits_{j=0}^{\infty} \widetilde{D}_x^j \Big(2\, \sqrt{|v_1|} - u_x\, v_1 \Big)\; \frac{\partial}{\partial v_j}, \end{array} \right. \tag{38}$$

$$\left\{ \begin{array}{lcl} \widetilde{D}_t & = & \bar{D}_t - \sum\limits_{j=0}^{\infty} \widetilde{D}_x^j\, (u + (u_x^2 + u_y)\, v_1)\; \frac{\partial}{\partial v_j}, \\ \widetilde{D}_x & = & \bar{D}_x + \sum\limits_{j=0}^{\infty} v_{j+1}\, \frac{\partial}{\partial v_j}, \\ \widetilde{D}_y & = & \bar{D}_y - \sum\limits_{j=0}^{\infty} \widetilde{D}_x^j\, (v_1\, u_x + x)\, \frac{\partial}{\partial v_j}, \end{array} \right. \tag{39}$$

$$\left\{\begin{array}{lcl} \widetilde{D}_t &=& \bar{D}_t - \sum\limits_{j=0}^{\infty} \widetilde{D}_x^j \left(\left(u_y + \lambda\, u_x - \lambda^2\right) v_1\right) \frac{\partial}{\partial v_j}, \\ \widetilde{D}_x &=& \bar{D}_x + \sum\limits_{j=0}^{\infty} v_{j+1} \frac{\partial}{\partial v_j}, \\ \widetilde{D}_y &=& \bar{D}_y - \sum\limits_{j=0}^{\infty} \widetilde{D}_x^j \left(\left(u_x - \lambda\right) v_1\right) \frac{\partial}{\partial v_j}. \end{array}\right. \tag{40}$$

The covering (36) was found in the cases of $\kappa = 0$ and $\kappa = 1$ in [9] and [21], respectively. The covering (40) was found in [11]. In [36], we construct the coverings (35), (36) for $\kappa \not\in \{-2, -\frac{3}{2}, -1\}$, and the covering (40) by means of another technique. Another covering of Eq. (10) in the case of $\kappa = -1$ was found in [36], too. As far as we know, the coverings (37), (38), and (39) are new.

5. Conclusion

We have shown that the approach based on the contact integrable extensions allows one to derive coverings of PDEs with three independent variables from MC forms of their symmetry pseudo-groups. We hope that this technique will be effective in studying a wide variety of PDEs with large symmetry pseudo-groups. Two important problems require further exploration. The first one is a connection between the restrictions like (20) for the form of a CIE and the number of the additional invariants with structure properties of the symmetry pseudo-groups. The second task is a refinement of the analysis of over-determined systems of PDEs for the coefficients of CIEs by means of Cartan's method. We hope to address these problems in future papers.

Bibliography

[1] Błaszak, M.: Classical R-matrices on Poisson algebras and related dispersionless systems, Phys. Lett. A **297**, 191–195 (2002)

[2] Bogdanov, L.V., Konopelchenko, B.G.: Nonlinear Beltrami equations and τ-functions for dispersionless hierarchies, Phys. Lett. A **322**, 330–337 (2004)

[3] Bryant, R.L., Chern, S.S., Gardner, R.B., Goldschmidt, H.L., Griffiths, P.A.: Exterior Differential Systems. N.Y., Springer-Verlag (1991)

[4] Bryant, R.L., Griffiths, Ph.A.: Characteristic cohomology of differential systems (II): conservation laws for a class of parabolic equations, Duke Math. J. **78**, 531–676 (1995)

[5] Cartan, É.: Sur la structure des groupes infinis de transformations. Œuvres Complètes, Part II, **2**, p. 571–715. Gauthier - Villars, Paris (1953)

[6] Cartan, É.: Les sous-groupes des groupes continus de transformations. Œuvres Complètes, Part II, **2**, p. 719–856. Gauthier - Villars, Paris (1953)

[7] Cartan, É.: Les problèmes d'équivalence. Œuvres Complètes, Part II, **2**, p. 1311–1334. Gauthier - Villars, Paris (1953)

[8] Cartan, É.: La structure des groupes infinis. Œuvres Complètes, Part II, **2**, p. 1335–1384. Gauthier - Villars, Paris (1953)

[9] Chang, J.-H., Tu, M.-H.: On the Miura map between the dispersionless KP and dispersionless modified KP hierarchies. J. Math. Phys., **41**, 5391–5406 (2000).

[10] Dodd, R., Fordy, A.: The prolongation structures of quasipolynomial flows. Proc. Roy. Soc. London, A, **385**, 389–429 (1983)

[11] Dunajski, M.: A class of Einstein–Weil spaces associated to an integrable system of hydrodynamic type, J. Geom. Phys. **51**, 126-137 (2004)

[12] Estabrook, F.B.: Moving frames and prolongation algebras. J. Math. Phys., **23** (1982), 2071–2076 (1982)

[13] Fels, M., Olver, P.J.: Moving coframes. I. A practical algorithm. Acta. Appl. Math. **51**, 161–213 (1998)

[14] Ferapontov, E.V., Khusnutdinova, K.R.: The characterization of two-component (2+1)-dimensional integrable systems of hydrodynamic type, J. Phys. A.: Math. Gen. **37** 2949–2963 (2004)

[15] Gardner, R.B.: The method of equivalence and its applications. CBMS–NSF regional conference series in applied math., SIAM, Philadelphia (1989)

[16] Harrison, B.K.: On methods of finding Bäcklund transformations in systems with more than two independent variables. J. Nonlinear Math. Phys., **2**, 201–215 (1995)

[17] Harrison, B.K.: Matrix methods of searching for Lax pairs and a paper by Estévez. Proc. Inst. Math. NAS Ukraine, **30**, Part 1, 17–24 (2000)

[18] Hoenselaers, C.: More prolongation structures. Prog. Theor. Phys. **75**, 1014–1029 (1986)

[19] Igonin, S.: Coverings and the fundamental group for partial differential equations. J. Geom. Phys., **56**, 939–998 (2006).

[20] Kamran, N.: Contributions to the Study of the Equivalence Problem of Élie Cartan and its Applications to Partial and Ordinary Differential Equations. Mem. Cl. Sci. Acad. Roy. Belg., **45**, Fac. 7 (1989)

[21] Konopelchenko, B., Martínez Alonso, L.: Dispersionless scalar hierarchies, Whitham hierarchy and the quasi-classical $\bar{\partial}$-method, J. Math. Phys. **43** 3807–3823 (2003)

[22] Krasil'shchik, I.S., Vinogradov, A.M.: Nonlocal symmetries and the theory of coverings. Acta Appl. Math., **2**, 79–86 (1984)

[23] Krasil'shchik, I.S., Lychagin, V.V., Vinogradov, A.M.: Geometry of Jet Spaces and Nonlinear Partial Differential Equations. Gordon and Breach, New York (1986)

[24] Krasil'shchik, I.S., Vinogradov, A.M.: Nonlocal trends in the geometry of differential equations: symmetries, conservation laws, and Bäcklund transformations. Acta Appl. Math., **15**, 161–209 (1989)

[25] Krasil'shchik, I.S., Vinogradov, A.M. (eds.): Symmetries and Conservation Laws for Differential Equations of Mathematical Physics. Transl. Math. Monographs 182, Amer. Math. Soc., Providence (1999).

[26] Krichever, I.M.: The averaging method for two-dimensional "integrable" equations. Funct.Anal. Appl. **22**, 200–213 (1988)

[27] Kupershmidt, B.A.: The quasiclassical limit of the modified KP hierarchy. J. Phys. A Math. Gen. **23**, 871–886 (1990)

[28] Kuz'mina, G.M.: On a possibility to reduce a system of two first-order partial differential equations to a single equation of the second order. Proc. Moscow State Pedagog. Inst. **271**, 67–76 (1967) (in Russian)

[29] Marvan, M.: On zero-curvature representations of partial differential equations. Proc. Conf. on Diff. Geom. and Its Appl., Opava (Czech Republic), 103–122 (1992)

[30] Marvan, M.: A direct procedure to compute zero-curvature representations. The case $\mathfrak{sl}_2$. In: Proc. Int. Conf. on Secondary Calculus and Cohomological Physics, Moscow, Russia, August 24-31, 1997. Available via the Internet at ELibEMS, `http://www.emis.de/proceedings`.

[31] Marvan, M.: On the horizontal gauge cohomology and nonremovability of the spectral parameter. Acta Appl. Math **72**, 51–65 (2002)

[32] Morozov, O.I.: Moving coframes and symmetries of differential equations. J. Phys. A, Math. Gen., **35**, 2965–2977 (2002)

[33] Morozov, O.I.: Contact-equivalence problem for linear hyperbolic equations. J. Math. Sci., **135**, 2680–2694 (2006)

[34] Morozov, O.I.: Coverings of differential equations and Cartan's structure theory of Lie pseudo-

groups. Acta Appl. Math. **99**, 309–319 (2007)

[35] Morozov, O.I.: Cartan's structure theory of symmetry pseudo-groups, coverings and multi-valued solutions for the Khokhlov–Zabolotskaya equation, Acta Appl. Math. **101**, 231-241 (2008)

[36] Morozov, O.I.: Cartan's structure of symmetry pseudo-group and coverings for the r-th modified dispersionless Kadomtsev–Petviashvili equation. Preprint `arXiv:0803.096v1 [nlin.SI]` (2008)

[37] Morris, H.C.: Prolongation structures and nonlinear evolution equations in two spatial dimensions. J. Math. Phys., **17**, 1870–1872 (1976)

[38] Morris, H.C.: Prolongation structures and nonlinear evolution equations in two spatial dimensions: a general class of equations. J. Phys. A, Math. Gen., **12**, 261–267 (1979)

[39] Olver, P.J.: Equivalence, Invariants, and Symmetry. Cambridge, Cambridge University Press (1995)

[40] Sakovich, S.Yu.: On zero-curvature representations of evolution equations. J. Phys. A, Math. Gen., **28**, 2861–2869 (1995)

[41] Stormark, O.: Lie's Structural Approach to PDE Systems. Cambridge, Cambridge University Press (2000)

[42] Takasaki, K.: Quasi-classical limit of BKP hierarchy and W-infinity symmetries, Lett. Math. Phys. **28**, 177–185 (1993)

[43] Tondo, G.S.: The eigenvalue problem for the three-wave resonant interaction in (2+1) dimensions via the prolongation structure. Lett. Nuovo Cimento, **44**, 297–302 (1985)

[44] Vasil'eva, M.V.: The Structure of Infinite Lie Groups of Transformations. Moscow, MGPI (1972) (in Russian)

[45] Wahlquist, H.D., Estabrook F.B.: Prolongation structures of nonlinear evolution equations. J. Math. Phys., **16**, 1–7 (1975)

[46] Zakharov, V.E.: Integrable systems in multidimensional spaces. Lect. Notes Phys., **153**, 190–216 (1982)

Appendix

The structure equations of the symmetry pseudo-group of Eq. (10) in the case of $\kappa \neq -1$ read

$$
\begin{aligned}
d\theta_0 &= \eta_1 \wedge \theta_0 + \xi^1 \wedge \theta_1 + \xi^2 \wedge \theta_2 + \xi^3 \wedge \theta_3, \\
d\theta_1 &= \tfrac{1}{16}\,\big(24\,\eta_1 - 24\,\theta_{22} - 12\,V\,\xi^1 - 12\,\xi^2 - (12\,U - \kappa + 4)\,\xi^3\big) \wedge \theta_1 + \xi^1 \wedge \theta_{11} + \xi^2 \wedge \theta_{12} \\
&\quad + \big(\tfrac{1}{8}\,(2\,\kappa^2 + 15\,\kappa + 4)\,\theta_2 + \kappa\,\theta_{23} + (\kappa\,U + 1 - \tfrac{1}{8}\,\kappa\,(\kappa - 10))\,\xi^2 \\
&\quad + \tfrac{1}{16}\,(7\,\kappa + 4)\,V\,\xi^3\big) \wedge \theta_0 + ((\kappa + 1)\,\theta_2 + (\kappa + 2)\,\xi^2) \wedge \theta_3 + \xi^3 \wedge \theta_{13}, \\
d\theta_2 &= \tfrac{1}{2}\,\big(\eta_1 - \theta_{22} - \tfrac{1}{2}\,V\,\xi^1 - \xi^2 - \big(U - \tfrac{7}{8}\,\kappa - \tfrac{1}{2}\big)\,\xi^3\big) \wedge \theta_2 + \xi^1 \wedge \theta_{12} + \xi^2 \wedge \theta_{22} + \xi^3 \wedge \theta_{23}, \\
d\theta_3 &= \big(\eta_1 - \theta_{22} - \tfrac{1}{2}\,V\,\xi^1 - \xi^2 - \big(U - \tfrac{1}{4}\,\kappa\big)\,\xi^3\big) \wedge \theta_3 + \tfrac{1}{2}\,(\kappa + 2)\,\xi^2 \wedge \theta_2 + \xi^1 \wedge \theta_{13} \\
&\quad + \tfrac{1}{8}\,(\kappa - 4)\,\big(\theta_{22} + \xi^2 + \tfrac{1}{8}\,(8\,U - \kappa - 4)\,\xi^3\big) \wedge \theta_0 + \xi^2 \wedge \theta_{23} + \xi^3 \wedge \theta_{12}, \\
d\xi^1 &= -\tfrac{1}{2}\,\big(\eta_1 - 3\,\theta_{22} + 3\,\xi^2 - \big(3\,U - \tfrac{1}{8}\,(\kappa - 4)\big)\,\xi^3\big) \wedge \xi^1, \\
d\xi^2 &= \tfrac{1}{2}\,\big(\eta_1 + \theta_{22} + \tfrac{1}{2}\,V\,\xi^1 + \big(U + \tfrac{1}{8}\,\kappa + \tfrac{3}{2}\big)\,\xi^3\big) \wedge \xi^2 + \big(\tfrac{1}{8}\,(\kappa - 4)\,\theta_0 - \theta_3\big) \wedge \xi^1 - \theta_2 \wedge \xi^3, \\
d\xi^3 &= \big(\theta_{22} + \tfrac{1}{2}\,V\,\xi^1 + \xi^2\big) \wedge \xi^3 - (\kappa + 2)\,\big(\theta_2 + \xi^2\big) \wedge \xi^1, \\
d\theta_{11} &= \big(2\,\eta_1 - 3\,\theta_{22} - 3\,\xi^2 - \big(3\,U - \tfrac{1}{8}\,(\kappa - 4)\big)\,\xi^3\big) \wedge \theta_{11} - \xi^1 \wedge \eta_7 - \xi^2 \wedge \eta_5 - \xi^3 \wedge \eta_6 \\
&\quad - \big(\kappa\,\eta_4 - \tfrac{3}{16}\,V\,\big(3\,\kappa^2 + 11\,\kappa + 4\big)\,\theta_2 - \kappa\,(U + 1)\,\theta_3 - \tfrac{1}{4}\,\kappa\,(\kappa + 8)\,\theta_{12} - \tfrac{3}{4}\,\kappa\,V\,\theta_{23} \\
&\quad + \tfrac{1}{64}\,(\kappa + 6)\,(\kappa - 4)\,(7\,\kappa + 4)\,V\,\xi^2 - \tfrac{5}{64}\,(7\,\kappa + 4)\,V^2\,\xi^3\big) \wedge \theta_0 + V\,\xi^2 \wedge \theta_{12} \\
&\quad + \big(\tfrac{1}{4}\,(5\,\kappa^2 + 31\,\kappa + 24)\,\theta_2 + (4\,\kappa + 3)\,\theta_{23} + \big((4\,\kappa + 3)\,U - \tfrac{1}{4}\,\kappa^2 + \tfrac{9}{2}\,\kappa + 5\big)\,\xi^2 \\
&\quad + \tfrac{1}{8}\,(15\,\kappa + 16)\,V\,\xi^3\big) \wedge \theta_1 + \big(\kappa\,\theta_{12} + \tfrac{1}{2}\,(\kappa + 2)\,V\,\xi^2\big) \wedge \theta_3 \\
&\quad + \big((2\,\kappa + 3)\,\theta_2 + 2\,(\kappa + 2)\,\xi^2 + \tfrac{5}{4}\,V\,\xi^3\big) \wedge \theta_{13},
\end{aligned}
\tag{41}
$$

$$
\begin{aligned}
d\theta_{12} =&\ \big(\eta_1 - 2\,\theta_{22} - 2\,\xi^2 - \big(2\,U - \tfrac{1}{2}\,\kappa\big)\ \xi^3\big) \wedge \theta_{12} - \xi^1 \wedge \eta_5 - \xi^2 \wedge \eta_3 - \xi^3 \wedge \eta_4 \\
&- \ \big(\theta_{23} + \big((\kappa+1)\,U + \tfrac{3}{8}\,\kappa^2 + \tfrac{9}{4}\,\kappa + 2\big)\,\xi^2 + \tfrac{1}{16}(11\,\kappa + 12)\,V\,\xi^3\big) \wedge \theta_2 \\
&- \ \big(\tfrac{1}{8}(\kappa - 4)\,\theta_0 + \theta_3 + \tfrac{1}{2}\,V\,\xi^2\big) \wedge \theta_{22} + \big((\kappa+2)\,\xi^2 + \tfrac{3}{4}\,V\,\xi^3\big) \wedge \theta_{23},
\end{aligned}
$$

$$
\begin{aligned}
d\theta_{13} =&\ \tfrac{1}{2}\ \big(3\,\eta_1 - 5\,\theta_{22} - 5\,\xi^2 - \big(5\,U - \tfrac{5}{8}\,\kappa + \tfrac{1}{2}\big)\ \xi^3\big) \wedge \theta_{13} - \xi^1 \wedge \eta_6 - \xi^2 \wedge \eta_4 - \xi^3 \wedge \eta_5 \\
&- \tfrac{1}{8}\,(\kappa - 4)\ \big(\eta_3 - (\kappa+2)\ \big(U - \tfrac{1}{8}\,\kappa\big)\ \theta_2 - \theta_3 - \tfrac{1}{8}\,V\,\theta_{22} - \theta_{23} \\
&- \ \tfrac{1}{8}\ \big((\kappa+2)\,(8\,U - \kappa - 4) + 6\,V\big)\ \xi^2 + \tfrac{1}{64}\,(\kappa+6)\,(7\kappa+4)\,V\,\xi^3\big) \wedge \theta_0 \\
&+ \tfrac{1}{8}\,(\kappa - 4)\ \big(\theta_{22} + \xi^2 + \big(U - \tfrac{1}{8}\,(\kappa+4)\big)\,\xi^3\big) \wedge \theta_1 \\
&- \ \big(\big(\tfrac{7}{8}\,\kappa^2 + 4\,\kappa + 3\big)\ \theta_3 + (\kappa+2)\,\theta_{12} + \tfrac{1}{4}\,(\kappa+2)\,V\,\xi^2\big) \wedge \theta_2 \\
&+ \ \big((2\,\kappa+1)\,\theta_{23} + \big(2\,(\kappa+1)\,U + \tfrac{1}{8}\,\kappa^2 + \tfrac{11}{4}\,\kappa + 3\big)\,\xi^2 + \tfrac{1}{16}\,(17\,\kappa + 20)\,V\,\xi^3\big) \wedge \theta_3 \\
&+ \ \big(\tfrac{3}{2}\,(\kappa+2)\,\xi^2 + V\,\xi^3\big) \wedge \theta_{12} + \tfrac{3}{4}\,V\,\xi^2 \wedge \theta_{23},
\end{aligned}
$$

$$
d\theta_{22} = \eta_3 \wedge \xi^1 - \tfrac{1}{2}\ (\eta_1 + \theta_{22}) \wedge \xi^2 + \eta_2 \wedge \xi^3
$$

$$
\begin{aligned}
d\theta_{23} =&\ \tfrac{1}{2}\ \big(\eta_1 - 3\,\theta_{22} - 3\,\xi^2 - \big(3\,U - \tfrac{11}{8}\,\kappa - \tfrac{1}{4}\big)\,\xi^3\big) \wedge \theta_{23} - \xi^1 \wedge \eta_4 - \xi^2 \wedge \eta_2 - \xi^3 \wedge \eta_3 \\
&- \frac{1}{8}\,(\kappa - 4)\,\xi^3 \wedge \theta_0 - \tfrac{1}{8}\ \big((3\,\kappa+4)\,\xi^2 - \big((5\,\kappa - 4)\,U + \tfrac{15}{8}\,\kappa^2 + 12\,\kappa + 10\big)\,\xi^3\big) \wedge \theta_2 \\
&+ \xi^3 \wedge \theta_3 + \tfrac{1}{8}\ \big(3\,(\kappa+4)\,\theta_2 + (8\,U - 3\,\kappa + 4)\,\xi^2 + 4\,V\,\xi^3\big) \wedge \theta_{22},
\end{aligned}
$$

$$
\begin{aligned}
d\eta_1 \ =&\ \tfrac{1}{4}\ \big(\kappa^2 + 7\,\kappa + 4\big)\ \xi^1 \wedge \theta_2 + \tfrac{1}{8}\,(\kappa - 4)\,\xi^3 \wedge \theta_{22} + \kappa\,\xi^1 \wedge \theta_{23} + \tfrac{1}{16}\,(7\,\kappa + 4)\,V\,\xi^1 \wedge \xi^3 \\
&+ \ \big(\kappa\,U - \tfrac{1}{8}\,\kappa^2 + \tfrac{5}{4}\,\kappa + 4\big)\ \xi^1 \wedge \xi^2 - \tfrac{1}{8}\,(\kappa - 4)\,\xi^2 \wedge \xi^3,
\end{aligned}
$$

$$
\begin{aligned}
d\eta_2 \ =&\ -\tfrac{1}{8}\ \big(4\,\theta_2 + \big(2\,U + \tfrac{1}{4}\,\kappa + 3\big)\big)\,\xi^2 - 3\,V\,\xi^3\big) \wedge \eta_1 - \big(\theta_{22} + V\,\xi^1 + \tfrac{1}{2}\,\xi^2 \\
&- \ (3\,U + 2\,\kappa + 1)\,\xi^3\big) \wedge \eta_2 - \tfrac{1}{32}\ \big(\big(32\,U - 7\,\kappa^2 + 38\,\kappa + 8\big)\ \xi^1 + 96\xi^3\big) \wedge \eta_3 \\
&+ \tfrac{1}{16}\,(\kappa - 4)\,\theta_0 \wedge \big(3\,U + \tfrac{1}{16}\ \big(7\,\kappa^2 + 40\,\kappa + 16\big)\ \xi^1 - 3\,\xi^3\big) \\
&- \tfrac{1}{2}\,\theta_2 \wedge \big(\theta_{22} + \big(\tfrac{1}{128}\,(\kappa+2)\,\big(8\,(7\kappa^2 + 126\kappa + 120)\,U + 21\kappa^3 + 198\kappa^2 + 688\kappa + 160\big) \\
&+ \ \tfrac{3}{2}\,(\kappa+3)\,V\big)\ \xi^1 + \tfrac{3}{8}\ \big(8\,\kappa\,U + 3\,\kappa^2 + 26\,\kappa + 24\big)\ \xi^3\big) - \tfrac{1}{32}\,\theta_3 \wedge \big((48\,U + 7\,\kappa^2 + 40\,\kappa \\
&+ \ 16)\,\xi^1 - 48\,\xi^3\big) - \theta_{12} \wedge \xi^1 + \tfrac{1}{64}\,\theta_{22} \wedge \big((7\,\kappa^2 + 38\,\kappa - 8)\,V\,\xi^1 - (16\,U + 6\,\kappa + 8)\,\xi^2 \\
&- \ (64\,(2\,\kappa+1)\,U + 24\,V + 128\,(\kappa+1))\,\xi^3\big) - \tfrac{1}{32}\,\theta_{23} \wedge \big((\kappa+2)\,(64\,U + 7\,\kappa^2 + 38\,\kappa \\
&- \ 8)\,\xi^1 - 16\,(3\,\kappa+4)\,\xi^3\big) + \big(128\,(\kappa+2)\,U^2 + 24\,U\,V + 4\,(\kappa+2)\,(7\,\kappa^2 + 70\,\kappa + 40)\,U \\
&+ \ (7\,\kappa^2 + 91\,\kappa + 148)\,V + 2\,(\kappa+2)\,(7\,\kappa^2 + 70\,\kappa + 24)\big)\ \xi^1 \wedge \xi^2 + \tfrac{1}{2048}\,V\ \big(32\,(7\,\kappa^2 \\
&+ \ 166\,\kappa + 152)\,U + 2304\,V - 49\,\kappa^4 - 756\,\kappa^3 - 2324\,\kappa^2 + 5152\,\kappa + 2752\big)\ \xi^1 \wedge \xi^3 \\
&+ \tfrac{1}{8}\ \big(3\,V - 6\,U^2 + 2\,(19\,\kappa + 16)\,U + \tfrac{1}{32}\ \big(31\,\kappa^2 + 1144\,\kappa + 1200\big)\big)\ \xi^2 \wedge \xi^3,
\end{aligned}
$$

$$
\begin{aligned}
d\eta_3 \ =&\ -\tfrac{1}{16}\ \big(8\,\eta_3 - (\kappa - 4)\,\theta_0 + 8\,\theta_3 + 2\,V\,\xi^2\big) \wedge \eta_1 + \big((\kappa+2)\,(\theta_2 + \xi^2) - \tfrac{1}{2}\,V\,\xi^3\big) \wedge \eta_2 \\
&- \tfrac{1}{32}\ \big(48\,\theta_{22} - 24\,V\,\xi^1 + 32\,\xi^2 + (80\,U + \kappa\,(7\,\kappa + 36))\,\xi^3\big) \wedge \eta_3 + (\kappa+2)\,\eta_4 \wedge \xi^1 \\
&+ \tfrac{1}{16}\,(\kappa - 4)\,\theta_0 \wedge \big(\theta_{22} + (2\,(\kappa+2)\,(2\,U + 1) - V)\ \xi^1 + \tfrac{1}{16}\,(48\,U + 7\,\kappa^2 + 40\,\kappa + 16)\xi^3\big) \\
&+ \tfrac{1}{8}\,(\kappa - 4)\,\theta_1 \wedge \xi^1 + \tfrac{1}{32}\,\theta_2 \wedge \big(V\,(80\,(\kappa+2)\,U + 7\,\kappa^3 + 36\,\kappa^2 + 10\,\kappa - 56)\,\xi^1 \\
&+ 4\,(\kappa^2 + 7\,\kappa + 4)\,\xi^2 - \big((\kappa+2)\,(7\,\kappa^2 + 126\,\kappa + 120)\,U + 24\,(\kappa+3)\,V \\
&+ \ \tfrac{1}{8}\,(\kappa+2)\,(21\,\kappa^3 + 198\,\kappa^2 + 688\,\kappa + 160)\big)\ \xi^3\big) - \tfrac{1}{2}\,\theta_3 \wedge (\theta_{22} + 2\,((\kappa+2)\,(2\,U + 1) \\
&- \ V)\,\xi^1 + \big(6\,U + \tfrac{7}{16}\,\kappa^2 + \tfrac{5}{2}\,\kappa + 1\big)\,\xi^3\big) + \big((\kappa+2)\ \big(U + \tfrac{3}{8}\,(\kappa+4)\big)\ \xi^1 + \xi^3\big) \wedge \theta_{12} \\
&- \theta_{13} \wedge \xi^1 + \tfrac{1}{64}\,V\,\theta_{22} \wedge \big(8\,\xi^2 - \big(7\,\kappa^2 + 38\,\kappa - 8\big)\ \xi^3\big) - \tfrac{1}{4}\,\theta_{23} \wedge \big(\kappa\,(V\,\xi^1 - 2\,\xi^2) \\
&+ \ \tfrac{1}{8}\,(\kappa+2)\,(64\,U + 7\,\kappa^2 + 38\,\kappa - 8)\,\xi^3\big) - \tfrac{1}{64}\,V\ \big(32\,(11\,\kappa + 5)\,U + 8\,V + 21\,\kappa^3 \\
&+ \ 124\,\kappa^2 + 44\,\kappa - 208\big)\ \xi^1 \wedge \xi^2 - \tfrac{1}{256}\,V^2\ \big(160\,U + 21\,\kappa^2 + 210\,\kappa + 104\big)\,\xi^1 \wedge \xi^3 \\
&- \tfrac{1}{64}\ \big(128\,(\kappa+2)\,U^2 + 24\,U\,V + 4\,(\kappa+2)\,(7\,\kappa^2 + 70\,\kappa + 40)\,U + (7\,\kappa^2 + 105\,\kappa + 156)\,V \\
&+ \ 2\,(\kappa+2)\,(7\,\kappa^2 + 70\,\kappa + 24)\big)\,\xi^2 \wedge \xi^3,
\end{aligned}
$$

$$
\begin{aligned}
d\eta_4 \ &= \eta_8 \wedge \xi^1 + \big(\theta_3 - \tfrac{1}{8}\,(\kappa-4)\,\theta_0 - \tfrac{3}{4}\,V\,\xi^2\big) \wedge \eta_2 + \tfrac{1}{32}\,\,(8\,(11\,\kappa+7)\,\theta_2 + 48\,\theta_{23} \\
&-\ \ (7\,\kappa^2 + 18\,\kappa - 88)\,\xi^2 + 56\,\xi^3\big) \wedge \eta_3 + \big(\eta_1 - 3\,\theta_{22} - 3\,\xi^2 - \big(3\,U - \tfrac{1}{4}\,\kappa + 2\big)\,\,\xi^3\big) \wedge \eta_4 \\
&+\ \tfrac{1}{256}\,(\kappa-4)\,\,\big(4\,(11\,\kappa+20)\,\theta_2 - 4\,(8\,U - 3\,\kappa + 4)\,\theta_{22} + 48\,\theta_{23} - (48\,U + 7\,\kappa^2 + 8\,\kappa - 48)\,\xi^2 \\
&-\ \ 32\,((\kappa+2)\,(2\,U+1) - V)\,\xi^3\big) \wedge \theta_0 + \tfrac{1}{512}\,\,(64\,(11\,\kappa+20)\,\theta_3 + 16\,V\,(13\,\kappa+20)\,\theta_{22} \\
&-\ \ 32\,(24\,(\kappa+2)\,U - 13\,\kappa^2 - 40\,\kappa - 16)\,\theta_{23} + 2\,\,\big(8\,(\kappa+2)\,(7\,\kappa^2 + 106\,\kappa + 136)\,U \\
&+\ \ 192\,(\kappa+3)\,V + (\kappa+2)\,(21\,\kappa^3 + 138\,\kappa^2 + 304\,\kappa - 160)\big)\,\,\xi^2 - V\,\,(16\,(65\,\kappa+172)\,U \\
&+\ \ 77\,\kappa^3 + 444\,\kappa^2 + 544\,\kappa - 768\big)\,\,\xi^3\big) \wedge \theta_2 + \big(\big(U - \tfrac{1}{8}\,(3\,\kappa-4)\big)\,\,\theta_{22} - \tfrac{3}{2}\,\theta_{23} \\
&+\ \ \tfrac{1}{32}\,\,\big(48\,U + 7\,\kappa^2 + 8\,\kappa - 48\big)\,\,\xi^2 + ((\kappa+2)\,(2\,U+1) - V)\,\xi^3\big) \wedge \theta_3 \\
&+\ \tfrac{1}{64}\,\,\big(24\,(\kappa+4)\,(\theta_{22} + \xi^2) + (8\,(3\,\kappa+20)\,U + 9\,\kappa^2 + 48\,\kappa + 112)\,\xi^3\big) \wedge \theta_{12} \\
&+\ \tfrac{1}{32}\,V\,\,\big((24\,U + 7\,\kappa^2 + 25\,\kappa - 44)\,\xi^2 - 8\,V\,\xi^3\big) \wedge \theta_{22} + \tfrac{1}{128}\,\,(4\,(32\,(3\,\kappa+7)\,U + 12\,V \\
&+\ \ 7\,\kappa^3 + 32\,\kappa^2 - 4\,\kappa - 48)\,\xi^2 - V\,(96\,U + 21\,\kappa^2 + 154\,\kappa + 8)\,\xi^3\big) \wedge \theta_{23} \\
&+\ \tfrac{1}{2048}\,V\,\,\big(32\,(7\,\kappa^2 + 236\,\kappa + 504)\,U + 2560\,V - 49\,\kappa^4 - 84\,\kappa^3 + 1644\,\kappa^2 + 6560\,\kappa \\
&-\ \ 3904)\,\,\xi^2 \wedge \xi^3, \\
d\eta_5 \ &= \eta_9 \wedge \xi^1 + \eta_8 \wedge \xi^3 - \big(\tfrac{1}{4}\,(\kappa-4)\,\theta_0 - 2\,\theta_3 - 2\,\theta_{12} - \tfrac{3}{2}\,V\,\xi^2\big) \wedge \eta_3 + ((\kappa+3)\,\theta_2 \\
&+\ \ (\kappa+2)\,\xi^2 + \tfrac{5}{4}\,V\,\xi^3\big) \wedge \eta_4 + \tfrac{1}{16}\,\,\big(24\,\eta_1 - 56\,(\theta_{22} + \xi^2) - (56\,U - 9\,\kappa + 8)\,\xi^3\big) \wedge \eta_5 \\
&+\ \tfrac{1}{64}\,(\kappa-4)\,\theta_0 \wedge \big((8\,(\kappa+1)\,U + 3\,\kappa^2 + 18\,\kappa + 16)\,\theta_2 - 16\,\theta_{12} + 4\,V\,\theta_{22} + 8\,(\kappa+2)\,\theta_{23} \\
&+\ \ 4\,(2\,(\kappa+2)\,(2\,U+1) - V)\,\xi^2\big) + \tfrac{1}{8}\,(\kappa-4)\,\theta_1 \wedge (\theta_{22} + \xi^2) \\
&+\ \tfrac{1}{64}\,\theta_2 \wedge \big(8\,\,\big(8\,(\kappa+1)\,U + 3\,\kappa^2 + 18\,\kappa - 16\big)\,\,\theta_3 + 16\,\,\big(8\,(\kappa+2) - 3\,\kappa^2 - 11\,\kappa - 4\big)\,\,\theta_{12} \\
&-\ \ 16\,V\,(3\,\kappa+5)\,\theta_{23} + V\,\,\big(32\,(3\,\kappa+8)\,U + 7\,\kappa^3 + 71\,\kappa^2 + 126\,\kappa - 8\big)\,\,\xi^2 \\
&+\ \ V^2\,(11\,\kappa+12)\,\xi^3\big) + \theta_3 \wedge \big(+2\,\theta_{12} - (\kappa+2)\theta_{23} - ((\kappa+2)\,(2\,U+1) - V)\,\,\xi^2\big) \\
&+\ \theta_{12} \wedge \big((\kappa-1)\,\theta_{23} - \big((\kappa+5)\,U + \tfrac{1}{2}\,V - \tfrac{3}{8}\,\kappa\,(\kappa+6)\big)\,\xi^2 - \tfrac{1}{32}\,V\,\,\big(32\,U + 7\,\kappa^2 + 82\,\kappa \\
&+\ \ 24)\,\,\xi^3\big) - \theta_{13} \wedge (\theta_{22} + \xi^2) - \tfrac{1}{8}\,V^2\,\theta_{22} \wedge \xi^2 + \tfrac{1}{8}\,V\,\theta_{23} \wedge \big(2, (3\,\kappa+8)\,\xi^2 + 3\,V\,\xi^3\big) \\
&+\ \tfrac{1}{256}\,V^2\,\big(160\,U + 21\,\kappa^2 + 210\,\kappa + 104\big)\,\,\xi^2 \wedge \xi^3, \\
d\eta_6 \ &= \eta_{10} \wedge \xi^1 + \eta_8 \wedge \xi^2 + \eta_9 \wedge \xi^3 + \tfrac{1}{32}\,\eta_3 \wedge ((\kappa-4)\,(5\,V\,\theta_0 + 8\,\theta_1) - 40\,\theta_{13}) \\
&+\ \eta_4 \wedge \big(2\,\kappa\,\theta_3 - \tfrac{1}{8}\,\kappa\,(\kappa-4)\,\theta_0 + V\,\xi^2\big) - \tfrac{1}{2}\,\eta_5 \wedge \big((\kappa+2)\,(4\,\theta_2 + 5\,\xi^2) + 3\,V\,\xi^3\big) \\
&-\ \eta_6 \wedge \big(\ 2\,(\eta_1 + 2\,\theta_{22} + 2\,\xi^2) + \big(4\,U - \tfrac{1}{8}\,(3\,\kappa-4)\big)\,\,\xi^3\big) + \tfrac{1}{2048}\,(\kappa-4)\,\theta_0 \wedge (64\,(\kappa-4)\,\theta_1 \\
&+\ \ 8\,(40\,(\kappa+2)\,U - 19\,\kappa^2 - 72\,\kappa - 56)\,\theta_2 + 128\,(\kappa+2)\,\theta_{12} + 640\,\theta_{13} - 64\,V\,\theta_{23} \\
&+\ \ 4\,V\,(16\,(7\,\kappa+16)\,U + 20\,V + 7\,\kappa^3 + 10\,\kappa^2 - 152\,\kappa - 256)\,\xi^2 - V^2\,(160\,U + 35\,\kappa^2 \\
&+\ \ 302\,\kappa + 152)\,\xi^3\big) + \tfrac{1}{32}\,\theta_1 \wedge \big(4\,(256\,\kappa^3 - 3\,\kappa - 4 - 2\,(\kappa-4)\,(\kappa+2)\,U)\,\theta_2 \\
&+\ \ (\kappa-4)\,(8\,\theta_3 - V\,\theta_{22} + 4\,\theta_{23} + ((\kappa+2)\,(8\,U - \kappa - 4) + V)\,\xi^2 + (40\,U + 7\,\kappa^2 \\
&+\ \ 41\,\kappa + 4)\,\xi^3)\big) - \theta_2 \wedge \big(\tfrac{1}{32}\,V\,(13\,\kappa^2 + 4\,\kappa - 8)\,\theta_3 + \tfrac{1}{16}\,(40\,(\kappa+2)\,U - 25\,\kappa^2 - 112\,\kappa \\
&-\ \ 64)\,\theta_{13}) - \tfrac{1}{32}\,\theta_3 \wedge (((\kappa-4)\,(4\,\kappa\,U + 5\,V) + 2\,\kappa\,(4\,\kappa-1))\,\theta_0 - 4\,\kappa\,(7\,\kappa+20)\,\theta_{12} \\
&+\ \ 8\,V\,(4\,\kappa+5)\,\theta_{23} + V\,(80\,(\kappa+1)\,U + 7\,\kappa^3 + 12\,\kappa^2 - 56\,\kappa - 64)\,\xi^2\big) \\
&+\ \tfrac{1}{64}\,(\kappa-4)\,\theta_{11} \wedge \big(8\,(\theta_{22} + \xi^2) + (8\,U - \kappa - 4)\,\xi^3\big) + \tfrac{1}{8}V\,\theta_{12} \wedge \big(13\,(\kappa+2)\,\xi^2 + 4\,V\,\xi^3\big) \\
&+\ \theta_{13} \wedge \big(\tfrac{1}{2}\,(7\,\kappa+2)\,\theta_{23} + \tfrac{1}{8}\,(8\,(\kappa-3)\,U - 5\,V + 8\,(5\,\kappa+4))\,\xi^2 + \tfrac{1}{128}\,V\,(160\,U + 35\,\kappa^2 \\
&+\ \ 430\,\kappa + 152)\,\xi^3\big) + \tfrac{3}{16}\,V^2\,\theta_{23} \wedge \xi^2, \\
d\eta_7 \ &= \eta_{11} \wedge \xi^1 + \eta_9 \wedge \xi^2 + \eta_{10} \wedge \xi^3 - \kappa\,\eta_8 \wedge \theta_0 - 3\,\eta_3 \wedge \theta_{11} + \eta_4 \wedge \big(\tfrac{3}{4}\,\kappa\,V\,\theta_0 + (5\,\kappa+3)\,\theta_1\big) \\
&+\ \tfrac{1}{8}\,\eta_5 \wedge \big((2\,\kappa^2 + 17\,\kappa - 4)\,\theta_0 + 8\,(\kappa-1)\,\theta_3 - 10\,V\,\xi^2\big) - \eta_6 \wedge \big((3\,\kappa+5)\,\theta_2 + 3\,(\kappa+2)\,\xi^2 \\
&+\ \ \tfrac{7}{4}\,V\,\xi^3\big) - \tfrac{1}{16}\,\eta_7 \wedge \big(40\,\eta_1 - 72\,(\theta_{22} + \xi^2) - (72\,U - 3\,(\kappa-4))\,\xi^3\big)
\end{aligned}
$$

$$\begin{aligned}
&+ \tfrac{1}{256}\,\theta_0 \wedge \big(8\,(\kappa-4)\,(4\,(4\,\kappa+3)\,U-\kappa^2+18\,\kappa-20)\,\theta_1+4\,V^2\,(26\,\kappa^2+57\,\kappa+28)\,\theta_2\\
&-\ 8\,V\,(48\,\kappa\,U+7\,\kappa^3+26\,\kappa^2-64\,\kappa-32)\,\theta_3-96\,(\kappa-4)\,\theta_{11}-16\,V\,(9\,\kappa^2+35\,\kappa+4)\,\theta_{12}\\
&+\ 64\,(4\,\kappa\,U+\kappa^2+2\,\kappa-8)\,\theta_{13}-(7\,\kappa+4)\,(\kappa^2-18\,\kappa-64)\,V^2\,\xi^2-10\,(7\,\kappa+4)\,V^3\,\xi^3\big)\\
&+ \tfrac{1}{64}\,\theta_1 \wedge (4\,(\kappa-1)\,(19\,\kappa+20)\,V\,\theta_2+32\,(2\,(5\,\kappa+3)\,U-(8\,\kappa+5)\,(\kappa-2))\,\theta_3\\
&+\ 16\,(6\,\kappa^2+39\,\kappa+24)\,\theta_{12}-144\,(\kappa+1)\,V\,\theta_{23}-V\,(96\,(4\,\kappa+3)\,U+35\,\kappa^3+63\,\kappa^2\\
&-\ 302\,\kappa-216)\,\xi^2+(5\,\kappa-12)\,V^2\,\xi^3\big)+\tfrac{1}{8}\,\theta_2 \wedge \big((24\,(\kappa+2)\,U-15\,\kappa^2-88\,\kappa-56)\,\theta_{11}\\
&-\ 2\,(3\,\kappa+7)\,V\,\theta_{13})+\tfrac{1}{8}\,\theta_3 \wedge \big(24\,\theta_{11}-(\kappa+2)\,(4\,V\,\theta_{12}+16\,\theta_{13}+V^2\,\xi^2)\big)\\
&+ \theta_{11} \wedge \big(3\,(\kappa+2)\,\theta_{23}+\tfrac{1}{8}\,(24\,(\kappa-1)\,U-6\,V-3\,\kappa^2+54\,\kappa+48)\,\xi^2+\tfrac{3}{64}\,(32\,U+7\,\kappa^2\\
&+\ 98\,\kappa+40)\,\xi^3\big)-\theta_{12} \wedge \big(3\,(\kappa+1)\,\theta_{13}-\tfrac{1}{4}\,V^2\,\xi^2\big)+\tfrac{1}{8}\,V\,\theta_{13} \wedge \big(18\,(\kappa+2)\,\xi^2+5\,V\,\xi^3\big)\,.
\end{aligned}$$

The structure equations of the symmetry pseudo-group of Eq. (10) in the case of $\kappa=-1$ have the form

$$\begin{aligned}
d\theta_0 &= \eta_1 \wedge \theta_0+\xi^1 \wedge \theta_1+\xi^2 \wedge \theta_2+\xi^3 \wedge \theta_3,\\
d\theta_1 &= \tfrac{3}{2}\,\eta_1 \wedge \theta_1+2\,\eta_2 \wedge \theta_3+\eta_3 \wedge \theta_0-\tfrac{1}{8}\,\big(12\,\theta_{22}+16\,U\,\xi^1+12\,\xi^2+7\,\xi^3\big) \wedge \theta_1+\xi^1 \wedge \theta_{11}\\
&\quad + \xi^2 \wedge \theta_{12}+\xi^3 \wedge \theta_{13},\\
d\theta_2 &= \tfrac{1}{8}\,\big(4\,(\eta_1-\theta_{22}-\xi^2)-3\,\xi^3\big) \wedge \theta_2+\xi^1 \wedge \theta_{12}+\xi^2 \wedge \theta_{22}+\xi^3 \wedge \theta_{23},\\
d\theta_3 &= \big(\eta_1-\theta_{22}-U\,\xi^1-\xi^2-\tfrac{5}{8}\,\xi^3\big) \wedge \theta_3+\eta_2 \wedge \theta_2-\tfrac{5}{8}\,(\theta_{22}+\xi^2) \wedge \theta_0+\xi^1 \wedge \theta_{13}\\
&\quad + \xi^2 \wedge \theta_{23}+\xi^3 \wedge \theta_{12},\\
d\xi^1 &= \tfrac{1}{8}\,\big(12\,\theta_{22}-4\,\eta_1+12\,\xi^2+7\,\xi^3\big) \wedge \xi^1,\\
d\xi^2 &= -\tfrac{1}{8}\,(5\,\theta_0+8\,\theta_3) \wedge \xi^1+\tfrac{1}{8}\,\big(4\,(\eta_1+\theta_{22})+3\,\xi^3\big) \wedge \xi^2-(\eta_2+\theta_2) \wedge \xi^3,\\
d\xi^3 &= -(2\,\eta_2+\theta_2) \wedge \xi^1+(\theta_{22}+U\,\xi^1+\xi^2) \wedge \xi^3,
\end{aligned}$$

$$\begin{aligned}
d\theta_{11} &= 2\,\eta_1 \wedge \theta_{11}+\eta_2 \wedge \big(\tfrac{1}{2}\,U\,\theta_0+4\,\theta_{13}\big)+6\,\eta_3 \wedge \theta_1+\eta_4 \wedge \theta_0+\eta_7 \wedge \xi^1+\eta_5 \wedge \xi^2+\eta_6 \wedge \xi^3\\
&\quad + \tfrac{1}{8}\,\big(30\,\theta_1-24\,U\,\theta_2-11\,\theta_3-14\,\theta_{12}-24\,U\,\theta_{23}-22\,U\,\xi^2+15\,U^2\,\xi^3\big) \wedge \theta_0\\
&\quad + \tfrac{5}{8}\,\big(10\,\theta_2+8\,\theta_{23}+7\,\xi^2-4\,U\,\xi^3\big) \wedge \theta_1+(2\,U\,\theta_3-\theta_{13}) \wedge \theta_2-(\theta_{12}+U\,\xi^2) \wedge \theta_3\\
&\quad - \big(3\,(\theta_{22}+\xi^2)+\tfrac{7}{4}\,\xi^3\big) \wedge \theta_{11}+2\,U\,\xi^2 \wedge \theta_{12}+3\,U\,\xi^3 \wedge \theta_{13},
\end{aligned}\tag{42}$$

$$\begin{aligned}
d\theta_{12} &= \eta_1 \wedge \theta_{12}+\eta_2 \wedge \big(2\,\theta_{23}+\tfrac{3}{2}\,\xi^2\big)+\eta_3 \wedge (\theta_2+\xi^2)+\eta_4 \wedge \xi^3+\eta_5 \wedge \xi^1-\tfrac{5}{8}\,\theta_0 \wedge (\theta_2-\theta_{22})\\
&\quad + \tfrac{1}{4}\,\theta_2 \wedge \big(7\,\xi^2+2\,U\,\xi^3\big)+\theta_3 \wedge (\theta_{22}+\xi^2)+\theta_{12} \wedge \big(2\,(\theta_{22}+\xi^2)+\tfrac{5}{4}\,\xi^3\big)\\
&\quad + \theta_{23} \wedge \big(2\,\xi^2-U\,\xi^3\big)+\tfrac{1}{4}\,U\,\xi^2 \wedge \xi^3,\\
d\theta_{13} &= \tfrac{3}{2}\,\eta_1 \wedge \theta_{13}+3\,\eta_2 \wedge \big(\tfrac{5}{16}\,\theta_0+\theta_{12}\big)+\eta_3 \wedge \big(\tfrac{5}{8}\,\theta_0+3\,\theta_3\big)+\eta_4 \wedge \xi^2+\eta_5 \wedge \xi^3+\eta_6 \wedge \xi^1\\
&\quad - \tfrac{5}{64}\,\theta_0 \wedge \big(17\,\theta_2+24\theta_3-8\,U\,\theta_{22}+8\,\theta_{23}-(8\,U-14)\xi^2-2\,U\,\xi^3\big)+\tfrac{5}{8}\,\theta_1 \wedge (\theta_{22}+\xi^2)\\
&\quad + \tfrac{1}{4}\,\theta_2 \wedge \big(13\,\theta_3+4\,\theta_{12}+2\,U\,\xi^2\big)-\tfrac{1}{8}\,\theta_3 \wedge \big(16\,\theta_{23}+21\,\xi^2+12\,U\,\xi^3\big)+2\,U\,\xi^3 \wedge \theta_{12}\\
&\quad + \tfrac{1}{2}\,\theta_{13} \wedge \big(5\,(\theta_{22}+\xi^2)+3\,\xi^3\big)+U\,\xi^2 \wedge \theta_{23},\\
d\theta_{22} &= -\tfrac{1}{2}\,\eta_1 \wedge \xi^2+\tfrac{3}{2}\,\eta_2 \wedge \xi^1+\eta_2 \wedge \xi^3+\eta_3 \wedge \xi^1+\theta_2 \wedge \big(\tfrac{21}{8}\,\xi^1+\xi^3\big)+\theta_3 \wedge \xi^1\\
&\quad - \tfrac{1}{8}\,\theta_{22} \wedge \big(8\,\xi^2+3\,\xi^3\big)+2\,\theta_{23} \wedge \xi^1-\tfrac{1}{4}\,\xi^1 \wedge (7\,\xi^2-U\,\xi^3),\\
d\theta_{23} &= \tfrac{1}{2}\,\eta_1 \wedge \theta_{23}+\eta_2 \wedge \big(\theta_{22}+\xi^2+\tfrac{3}{2}\,\xi^3\big)+\eta_3 \wedge \xi^3+\eta_4 \wedge \xi^1-\tfrac{5}{8}\,\theta_0 \wedge \xi^3\\
&\quad + \tfrac{1}{16}\,\theta_2 \wedge \big(72\,(\theta_{22}+\xi^2)+49\,\xi^3\big)-\tfrac{3}{8}\,\theta_{22}\,\big(4\,\theta_{23}+3\,\xi^2\big)+\tfrac{3}{2}\,\theta_{23} \wedge \big(\xi^2+2\,\xi^3\big)+\tfrac{77}{32}\,\xi^2 \wedge \xi^3,\\
d\eta_1 &= -\eta_3 \wedge \xi^1+\tfrac{5}{8}\,\big((\theta_0-\theta_2) \wedge \xi^1+(\theta_{22}+\xi^2) \wedge \xi^3\big)\,,\\
d\eta_2 &= -\tfrac{1}{4}\,\eta_2 \wedge \big(2\,\eta_1-2\,\theta_{22}-4\,U\,\xi^1-2\,\xi^2-\xi^3\big)-\eta_3 \wedge \xi^3-\tfrac{5}{16}\,\theta_0 \wedge (\xi_1-2\,\xi^3)\\
&\quad + \theta_2 \wedge \big(U\,\xi^1-\tfrac{13}{8}\,\xi^3\big)-\tfrac{1}{2}\,\theta_3 \wedge \xi^1+\tfrac{1}{2}\,\theta_{22} \wedge \xi^2-\theta_{23} \wedge \xi^3-\tfrac{1}{16}\,(8\,U\,\xi^1-21\,\xi^3) \wedge \xi^2,\\
d\eta_3 &= \tfrac{1}{16}\,\eta_1 \wedge (5\,\theta_0+8\,\eta_3)-\tfrac{1}{16}\,\eta_2 \wedge \big(20\,\theta_{22}+8\,U\,\xi^1+20\,\xi^2+15\,\xi^3\big)-\eta_4 \wedge \xi^1-\tfrac{5}{8}\,\theta_1 \wedge \xi^1
\end{aligned}$$

$$
\begin{aligned}
&+ \tfrac{1}{4}\,\eta_3 \wedge \big(6\,\theta_{22} + 8\,U\,\xi^1 + 6\,\xi^2 + \xi^3\big) - \tfrac{5}{32}\,\theta_0 \wedge \left(6\,\theta_{22} + (4\,U - 3)\,\xi^1 + \frac{15}{16}\,\xi^2 \wedge \theta_0 + \xi^3\right) \\
&+ \tfrac{1}{64}\,\theta_2 \wedge \big(192\,U\,\xi^1 - 40\,\xi^2 - 85\,\xi^3\big) + \tfrac{1}{8}\,\theta_3 \wedge \big(6\,\xi^1 - 5\,\xi^3\big) + \tfrac{9}{8}\,\theta_{12} \wedge \xi^1 \\
&+ \tfrac{5}{8}\,\theta_{22} \wedge (\xi^2 + U\,\xi^3) + \theta_{23} \wedge \big(3\,U\,\xi^1 - \tfrac{5}{8}\,\xi^3\big) - \tfrac{1}{8}\,U\,\xi^1 \wedge (22\,\xi^2 + 15\,U\,\xi^3) \\
&+ \tfrac{5}{32}\,(4\,U - 7)\xi^2 \wedge \xi^3, \\
d\eta_4 &= \eta_8 \wedge \xi^1 + \tfrac{1}{16}\,\eta_2 \wedge \big(+48\,\eta_3 - 30\,\theta_0 + 89\,\theta_2 + 16\,U\,\theta_{22} + 92\,\theta_{23} + 2\,(8\,U + 39)\,\xi^2 - 12\,U\,\xi^3\big) \\
&- \tfrac{1}{8}\,\eta_3 \wedge \big(17\,\theta_2 + 16\,\theta_{23} + 9\,\xi^2 - 8\,U\,\xi^3\big) - \eta_4 \wedge \big(\eta_1 - 3\,\theta_{22} - 3\,\xi^2 - \tfrac{23}{8}\,\xi^3\big) \\
&+ \tfrac{5}{64}\,\theta_0 \wedge \big(17\,\theta_2 - 13\,\theta_{22} + 16\,\theta_{23} - 4\,\xi^2 - \big(8\,U + \tfrac{77}{4}\big)\,\xi^3\big) + \tfrac{1}{64}\,\theta_2 \wedge (64\,(U\,\theta_{22} + \theta_{23}) \\
&+ \ (64\,U + 91)\xi^2 + 74\,U\,\xi^3\big) - \tfrac{1}{32}\,\theta_3 \wedge \big(26\,(\theta_{22} + \xi^2) + 77\,\xi^3\big) - \tfrac{1}{16}\,\theta_{12} \wedge \big(18\,(\theta_{22} + \xi^2) \\
&+ \ 31\xi^3\big) - \tfrac{1}{2}\,U\,\theta_{22} \wedge \xi^2 + \tfrac{3}{8}\,\theta_{23} \wedge \big(\xi^2 + U\,\xi^3\big) + \tfrac{13}{16}\,U\,\xi^2 \wedge \xi^3, \\
d\eta_5 &= \eta_9 \wedge \xi^1 + \eta_8 \wedge \xi^3 - \tfrac{1}{8}\,\eta_2 \wedge \big(15\,\theta_0 - 4\,U\,\theta_2 + 24\,\theta_3 + 4\,\theta_{12} - 12\,U\,\xi^2\big) \\
&- \tfrac{1}{4}\,\eta_3 \wedge \big(5\,\theta_0 + 8\,\theta_3 - 8\,U\,\theta_2 + 16\,\theta_{12} - 8\,U\,\xi^2\big) - \eta_4 \wedge \big(4\,\eta_2 + 2\,\theta_2 - \xi^2 + 2\,U\,\xi^3\big) \\
&- \tfrac{1}{8}\,\eta_5 \wedge \big(12\,\eta_1 - 28\,(\theta_{22} + \xi^2) - 17\,\xi^3\big) - \tfrac{5}{64}\,\theta_0 \wedge ((16\,U - 41)\theta_2 - 16\,\theta_3 - 32\,\theta_{12} \\
&- \ 24\,\theta_{23} + 2\,(8\,U - 7)\xi^2 + 4\,U\,\xi^3\big) - \tfrac{5}{8}\,\theta_1 \wedge (\theta_{22} + \xi^2) - \tfrac{1}{8}\,\theta_2 \wedge (41\,\theta_3 + 46\,\theta_{12} \\
&- \ 16\,U\,\xi^2 - 11\,U^2\,\xi^3\big) + \tfrac{1}{4}\,\theta_3 \wedge \big(4\,U\,\theta_{22} + 12\,\theta_{23} + (4\,U - 7)\,\xi^2 \wedge \theta_3 - 2\,U\,\xi^3\big) \\
&+ \tfrac{1}{8}\,\theta_{12} \wedge \big(32\,\theta_{23} + 23\,\xi^2 - 18\,U\,\xi^3\big) - \theta_{13} \wedge (\theta_{22} + \xi^2) + U\,\theta_{23} \wedge \big(2\,(\theta_2 + \xi^2) + U\,\xi^3\big) \\
&+ \tfrac{13}{8}\,U^2\,\xi^2 \wedge \xi^3, \\
d\eta_6 &= \eta_{10} \wedge \xi^1 + \eta_8 \wedge \xi^2 + \eta_9 \wedge \xi^3 - \tfrac{1}{8}\,\eta_2 \wedge (15\,\theta_1 - 12\,U\,\theta_3 + 8\,U\,\theta_{12} - 6\,\theta_{13}) \\
&+ \tfrac{1}{8}\,\eta_3 \wedge (5\,U\,\theta_0 - 10\,\theta_1 + 72\,U\,\theta_3 - 56\,\theta_{13}) - \tfrac{1}{8}\,\eta_4 \wedge \big(5\,\theta_0 + 16\,\theta_3 + 8\,U\,\xi^2\big) \\
&- \eta_5 \wedge \big(+5\,\eta_2 + 2\,\theta_2 + 3\,U\,\xi^3\big) - \eta_6 \wedge \big(2\,\eta_1 - 4\,\theta_{22} - 4\,\xi^2 - \tfrac{19}{8}\,\xi^3\big) \\
&+ \tfrac{5}{64}\,\theta_0 \wedge \big(+10\,\theta_1 - 12\,U\,\theta_2 - (72\,U - 11)\theta_3 - 4\,\theta_{12} + 56\,\theta_{13} - 16\,U\,\theta_{23} - 14\,U\,\xi^2 \\
&+ \ 11\,U^2\,\xi^3\big) + \tfrac{5}{32}\,\theta_1 \wedge \big(19\,\theta_2 + 4\,U\,\theta_{22} + 12\,\theta_{23} + 2\,(4\,U + 7)\,\xi^2 - 2\,U\,\xi^3\big) \\
&+ \tfrac{1}{4}\,\theta_2 \wedge (53\,U\,\theta_3 + 4\,U\,\theta_{12} - 37, \theta_{13}) - \tfrac{1}{8}\,\theta_3 \wedge \big(13\,\theta_{12} + 96\,U\,\theta_{23} + 87\,U\,\xi^2 + 45\,U^2\,\xi^3\big) \\
&- \tfrac{5}{8}\,\theta_{11} \wedge (\theta_{22} + \xi^2) - 2\,U\,\theta_{12} \wedge (\xi^2 - U\xi^3) + \theta_{13} \wedge \big(7\,\theta_{23} - \tfrac{29}{8}\,U\,\xi^3\big)\,, \\
d\eta_7 &= \eta_8 \wedge \theta_0 + \eta_{11} \wedge \xi^1 + \eta_9 \wedge \xi^2 + \eta_{10} \wedge \xi^3 + \eta_3 \wedge (24\,U\,\theta_1 - 11\,\theta_{11}) - \eta_4 \wedge (3\,U\,\theta_0 + 2\,\theta_1) \\
&- \tfrac{1}{4}\,\eta_2 \wedge \big(13\,U^2\,\theta_0 - 6\,U\,\theta_1 + 4\,\theta_{11} + 8\,U\,\theta_{13}\big) - \tfrac{1}{8}\,\eta_5 \wedge \big(19\,\theta_0 + 16\,(\theta_3 + U\,\xi^2)\big) \\
&- 2\,\eta_6 \wedge (3\,\eta_2 + 2\,\theta_2 + 4\,U\,\xi^3) - \tfrac{1}{8}\,\eta_7 \wedge \big(20\,\eta_1 - 36\,(\theta_{22} + \xi^2) - 21\,\xi^3\big) \\
&- \tfrac{1}{64}\,\theta_0 \wedge \big(5\,(192\,U - 1)\,\theta_1 - 88\,U^2\,\theta_2 - 32\,U\,\theta_3 - 440\,\theta_{11} - 272\,U\,\theta_{12} - 168\,\theta_{13} \\
&+ \ 16\,U^2\,\xi^2 - 120\,U^3\,\xi^3\big) - \tfrac{1}{8}\,\theta_1 \wedge \big(12\,\theta_3 + 18\,\theta_{12} + 248\,U\,\theta_{23} + 224\,U\,\xi^2 - 125\,U^2\,\xi^3\big) \\
&+ \theta_2 \wedge \big(36\,U\,\theta_1 - 2\,U^2\,\theta_3 - \tfrac{109}{8}\,\theta_{11} + 3\,U\,\theta_{13}\big) - \theta_3 \wedge (3\,U\,\theta_{12} + 2\,\theta_{13} - U^2\,\xi^2) \\
&+ \theta_{11} \wedge \big(11\,\theta_{23} + \tfrac{77}{8}\,\xi^2 - 5\,U\,\xi^3\big) - 3\,U\,\theta_{13} \wedge (\xi^2 - \,U\,\xi^3)
\end{aligned}
$$